

Supplementary Table SV Interaction of DHEAS with diagnostic group with respect to NEFA parameters.

Dependent variables	Independent variables	β coefficient \pm SE	P-value	R ²
NEFA ₀ (μ mol/l)	Diagnostic group \times DHEAS	0.001 \pm 0.001	0.894	0.1
	DHEAS	-0.001 \pm 0.001	0.959	
	Diagnostic group	-0.079 \pm 0.050	0.121	
g _s (mmol/l)	Diagnostic group \times DHEAS	0.001 \pm 0.001	0.045	8
	DHEAS	-0.001 \pm 0.001	0.126	
	Diagnostic group	0.113 \pm 0.058	0.057	
log K _C (%/min)	Diagnostic group \times DHEAS	-0.001 \pm 0.001	0.065	6.8
	DHEAS	0.001 \pm 0.001	0.148	
	Diagnostic group	-0.127 \pm 0.076	0.101	
Latency (τ) (min)	Diagnostic group \times DHEAS	-0.001 \pm 0.001	0.544	2.2
	DHEAS	-0.001 \pm 0.001	0.984	
	Diagnostic group	-0.011 \pm 0.066	0.86	
log Φ (mmol/l)	Diagnostic group \times DHEAS	-0.001 \pm 0.001	0.327	2.8
	DHEAS	0.001 \pm 0.001	0.684	
	Diagnostic group	-0.214 \pm 0.130	0.107	
S _{FFA} (μ mol/l/min)	Diagnostic group \times DHEAS	0.003 \pm 0.001	0.243	5.1
	DHEAS	-0.001 \pm 0.001	0.78	
	Diagnostic group	0.028 \pm 0.066	0.675	
K _{FFA} (%/min)	Diagnostic group \times DHEAS	-0.002 \pm 0.001	0.466	1.1
	DHEAS	0.001 \pm 0.001	0.511	
	Diagnostic group	0.027 \pm 0.060	0.658	