

Psychometric Properties of the STAI-Y scales and Normative Data in an Italian Elderly Population [Aging Clinical and Experimental Research]

Ciro Rosario Ilardi^a, Nadia Gamboz^b, Alessandro Iavarone^{c*}, Sergio Chieffi^d, Maria Antonella Brandimonte^b

^aDepartment of Psychology, University of Campania “Luigi Vanvitelli”, Caserta, Italy

^bLaboratory of Experimental Psychology, Suor Orsola Benincasa University, Naples, Italy

^cNeurological Unit, CTO Hospital, AORN “Ospedali dei Colli”, Naples, Italy

^dDepartment of Experimental Medicine, University of Campania “Luigi Vanvitelli”, Naples, Italy

*Corresponding Author: Alessandro Iavarone

e-mail address: aleiavarone@gmail.com

ORCID: 0000-0002-9596-5811

Online Resource 2 Results of Crawford & Garthwaite’s test for raw scores n SDs below or above the normative sample mean, and for cutoff, in the S-Anxiety and T-Anxiety scales

Z plus 95% confidence limits					
Standard deviation	Raw score	z	Lower	Upper	Width
S-Anxiety Scale					
1 SD below	25	-1.054	-1.182	-0.925	0.25
1.5 SD below	20	-1.490	-1.640	-1.340	0.30
1 SD above	49	1.04	0.912	1.168	0.25
1.5 SD above	54	1.476	1.327	1.625	0.29
Above the cutoff	68	2.698	2.475	2.920	0.44
T-Anxiety Scale					
1 SD below	25	-1.060	-1.188	-0.931	0.25
1.5 SD below	22	-1.321	-1.462	-1.179	0.28
1 SD above	49	1.027	0.899	1.154	0.25
1.5 SD above	55	1.549	1.395	1.701	0.30
Above the cutoff	67	2.592	2.376	2.807	0.43

S-Anxiety Scale; $N = 361$, mean = 37.08, SD = 11.46, range = 20–76

T-Anxiety Scale; $N = 361$, mean = 37.19, SD = 11.50, range = 22–84