

On-line Table: Clinical and angiographic findings in 40 patients with MCA aneurysms treated with the pCONus device

Pt	Sex, Age (yr)	Dome/Neck (mm)	Angiographic Results				Modified Rankin Scale		
			Immediate	3–6 Months	1 Year	Retreatment	Initial	At Discharge	At 30 Days
1	F, 76	9.8:5.5	NR	C	C		0	1	1
2	F, 54	7:5.2	AR	NR	NR		0	0	0
3	F, 65	7.9:2.7	AR	C	C		0	0	0
4	M, 51	9.1:5.3	NR	NR	C		0	0	0
5	M, 46	11.1:10.6	AR	AR	AR(w)	X	3	2	1
6	M, 44	7.1:4	NR	AR		X	2	2	2
7	M, 51	14.3:13.8	AR	AR(w)		X	0	0	0
8	M, 62	5.1:6.6	NR	AR		X	0	0	0
9	F, 58	4.3:4.7	AR	C	C		0	0	0
10	F, 68	17.4:6.1	NR	NR	NR		0	0	0
11	F, 57	13.1:6.1	AR	AR(w)	AR(w)	X	0	0	0
12	F, 56	5.7:4.8	NR	C			0	0	0
13	F, 56	7.7:5.1	AR	NR			0	0	0
14	F, 53	11.1:8.5	NR	AR	AR(w)		2	2	1
15	M, 76	3.2:4.8	AR	C			3	3	0
16	F, 53	4.6:4.5	NR	C			0	0	0
17	F, 58	11.2:3.3	NR	C			0	0	0
18	F, 75	4.1:6.1	NR	NR			0	0	0
19	F, 68	2.7:5.3	AR	NR			0	0	0
20	M, 60	7.4:3.2	NR	C			0	0	0
21	F, 64	4.9:2.8	NR	C			0	0	0
22	F, 62	6.1:4.5	C	C			0	0	0
23	M, 61	7.6:5.2	NR	NR			0	0	0
24	F, 68	11.3:9.8	NR	NR			0	0	0
25	F, 63	4.6:4.9	C	C			0	0	0
26	F, 62	4.6:5	C				0	0	0
27	F, 56	6.3:5.5	C				0	0	0
28	F, 53	5.6:4.6	AR				0	0	0
29	M, 77	10.8:8.2	AR	AR(w)		X	0	0	0
30	F, 52	8:3.5	NR				1	1	1
31	F, 49	8.8:5.4	NR	C	C		0	0	0
32	M, 36	6.3:5.1	C	C	C		0	0	0
33	F, 50	15:9.2	NR	NR	NR		1	1	1
34	M, 49	6.8:6.4	C				1	1	1
35	F, 60	6.6:4.7	C				1	1	1
36	M, 41	4.2:5.1	NR	NR			0	0	0
37	F, 53	5.5:3.3	C	C	C		0	0	0
38	F, 53	7:5	C	C			0	0	0
39	F, 69	6:5	NR	NR		X	0	0	0
40	M, 50	7:5	C	C	C		0	0	0

Note:—Pt indicates patient; C, complete aneurysm occlusion; NR, neck remnant; AR, aneurysm remnant; w, worsening; X, retreatment was done.