

Associations between seven-year C-reactive protein trajectory or pack-years smoked with choroidal or retinal thicknesses in young adults

Authors: Samantha Sze-Yee Lee¹; Darren John Beales²; Fred K Chen^{1,3}; Seyhan Yazar^{1,4}; David Alonso-Caneiro⁵; David A Mackey^{1,6,7}

¹Centre for Ophthalmology and Visual Science (incorporating Lions Eye Institute,), University of Western Australia, Western Australia, Australia

²School of Physiotherapy and Exercise Science, Curtin University, Western Australia, Australia

³Department of Ophthalmology, Royal Perth Hospital, Western Australia, Australia

⁴Single Cell and Computational Genomics Lab, Garvan Institute of Medical Research, New South Wales, Australia

⁵Queensland University of Technology (QUT), Contact Lens and Visual Optics Laboratory, Centre for Vision and Eye Research, School of Optometry and Vision Science, Queensland, Australia

⁶Centre for Eye Research Australia, University of Melbourne, Royal Victorian Eye and Ear Hospital, Victoria, Australia

⁷School of Medicine, Menzies Research Institute Tasmania, University of Tasmania, Tasmania, Australia

Supplementary figure S1. Box plots of C-reactive protein at each follow-up according to trajectory class

