



**Supplementary Table S5. Overall survival in patients with liver impairment (ITT population)**

	Normal	Liver Impairment Subgroup		
	Group A (n=546)	Group B (n=294)	Group C (n=443)	Group D (n=34)
<b>Median OS, months (95% CI)</b>	17.5 (15.9–18.8)	13.2 (11.7–14.5)	12.3 (11.1–13.6)	12.3 (8.0–16.1)
<b>OS rate (95% CI)<sup>a</sup></b>				
<b>1 year</b>	0.672 (0.633–0.712)	0.542 (0.485–0.599)	0.505 (0.458–0.552)	0.529 (0.362–0.739)
<b>2 years</b>	0.370 (0.328–0.412)	0.199 (0.151–0.248)	0.183 (0.145–0.221)	0.206 (0.070–0.342)
<b>3 years</b>	0.212 (0.170–0.253)	0.098 (0.055–0.141)	0.085 (0.051–0.118)	0 (NE–NE)



<sup>a</sup>OS rate at 1, 2, and 3 years and the 95% CIs were calculated using Kaplan–Meier estimate and Greenwood’s formula.

Group A, no liver impairment; group B, increased AST and/or ALT only; group C, any abnormality except increased bilirubin; group D, increased bilirubin.

ALT, alanine aminotransferase; AST, aspartate aminotransferase; CI, confidence interval; ITT, intent to treat; NE, nonestimable; OS, overall survival.