CERTIFICATION REGARDING CONFLICTS OF INTEREST AND CONFIDENTIALITY OF INFORMATION

Each Covered Person agrees to comply with the provisions of these Policies so long as he/she is a Covered Person. By signing, you are confirming that you have read and understand the above Policy on Conflicts of Interest and Confidentiality and agree to abide by same during all times that you are a Covered Person, as defined in the Policy.

asy Basueto, Pho MPH. ABPP Casey B. Azuero, PhD, MPH, ABPP

Charles H. Bombardier, PhD, ABPP

Jesse R. Fann, MD, MPH

Monteorden

Donald D. Kautz, RN, PhD

John hott Ruhard

Cand David AU PLD

J. Scott Richards, PhD, ABPP

Sunil Sabharwal, MD

CERTIFICATION REGARDING CONSORTIUM POLICIES AND PROCEDURES

Each Covered Person agrees to comply with the provisions of the policies and procedures outlined in the Clinical Practice Guideline Orientation Manual so long as he/she is a Covered Person. By signing, you are confirming that you have read and understand the Clinical Practice Guidelines Orientation Manual Policies and Procedures and agree to abide by same during all times that you are a Covered Person.

Casey Baguero, PhD. MPH. ABPP

Casey B. Azuero, PhD, MPH, ABPP

Charles H. Bombardier, PhD, ABPP

Jesse R. Fann, MD, MPH

Monteorder

COURSE PUP PLD

Donald D. Kautz, RN, PhD

John fult Ruhard

J. Scott Richards, PhD, ABPP

Sunil Sabharwal, MD

Appendix A: Search Strategies

Ovid MEDLINE® <1946 to August Week 2 2016>

- 1. exp Spinal Cord Injuries/
- 2. spinal cord injur*.mp.
- 3. (spine or spinal) and sci).mp
- 4. or/1-3
- 5. exp Mental Disorders/
- 6. (mental adj3 (health or disorder*)).mp
- 7. (anxiety or depression or post-traumatic or posttrau matic or PTSD or bipolar or suicid* or "self-harm").ti,ab.
- 8. exp Substance-Related Disorders/
- 9. (drug* or alcohol* or illicit or substance) adj3 (use* or abus*)).ti,ab.
- 10. or/5-9
- 11. 4 and 10
- 12. limit 11 to English language
- 13. limit 11 to abstracts
- 14. 12 or 13
- 15. limit 14 to humans

Ovid MEDLINE® In-Process & Other Non-Indexed Citations <August 22, 2016>

- 1. spinal cord injur*.mp.
- 2. ((spine or spinal) and sci).mp
- 3. (mental adj3 (health or disorder*)).mp
- 4. (anxiety or depression or post-traumatic or posttraumatic or PTSD or bipolar or suicid* or "self-harm").ti,ab.
- 5. ((drug* or alcohol* or illicit or substance) adj3 (use* or abus*)).ti,ab.
- 6. (1 or 2) and (3 or 4 or 5)
- 7. limit 6 to English language

PsycINFO <1806 to July Week 4 2016>

- 1. exp Spinal Cord Injuries/
- 2. spinal cord injur*.mp.
- 3. ((spine or spinal) and sci).mp
- 4. or/1-3
- 5. exp Mental Disorders/
- 6. (mental adj3 (health or disorder*)).mp
- (anxiety or depression or post-traumatic or posttraumatic or PTSD or bipolar or suicid* or "self-harm").ti,ab.
- 8. exp "SUBSTANCE USE DISORDER"/
- 9. ((drug* or alcohol* or illicit or substance) adj3 (use* or abus*)).ti,ab.
- 10. or/5-8
- 11. 4 and 10

Cochrane Central Register of Controlled Trials <July 2016>

- 1. exp Spinal Cord Injuries/
- 2. spinal cord injur*.mp.
- 3. ((spine or spinal) and sci).mp

- 4. or 1-3
- 5. exp Mental Disorders/
- 6. (mental adj3 (health or disorder*)).mp
- (anxiety or depression or post-traumatic or posttraumatic or PTSD or bipolar or suicid* or "self-harm").ti,ab.
- 8. exp Substance-Related Disorders/
- 9. ((drug* or alcohol* or illicit or substance) adj3 (use* or abus*)).ti,ab.
- 10. or $\sqrt{5-9}$
- 11. 4 and 10

Cochrane Database of Systematic Reviews

<2005 to August 17, 2016>

1. (((spine or spinal) and cord and injur*) or sci).ti.

Appendix B. Included Studies

- 1. Abdul-Sattar AB. Predictors of functional outcome in patients with traumatic spinal cord injury after inpatient rehabilitation: in Saudi Arabia. *Neurorehabilitation*. 2014;35(2):341-347.
- 2. Ahoniemi E, Pohjolainen T, Kautiainen H. Survival after spinal cord injury in Finland. Journal of Rehabilitation Medicine. 2011;43(6):481-485.
- 3. Alschuler KN, Jensen MP, Sullivan-Singh SJ, Borson S, Smith AE, Molton IR. The association of age, pain, and fatigue with physical functioning and depressive symptoms in persons with spinal cord injury. Journal of Spinal Cord Medicine. 2013;36(5):483-491.
- 4. Arango-Lasprilla JC, Ketchum JM, Starkweather A, Nicholls E, Wilk AR. Factors predicting depression among persons with spinal cord injury 1 to 5 years post injury. Neurorehabilitation. 2011;29(1):9-21.
- 5. Bailey J, Dijkers MP, Gassaway J, et al. Relationship of nursing education and care management inpatient rehabilitation interventions and patient characteristics to outcomes following spinal cord injury: the SCIRehab project. Journal of Spinal Cord Medicine. 2012;35(6):593-610.
- 6. Banerjea R, Findley PA, Smith B, Findley T, Sambamoorthi U. Co-occurring medical and mental illness and substance use disorders among veteran clinic users with spinal cord injury patients with complexities. Spinal Cord. 2009;47(11):789-795.
- 7. Binks TM, Radnitz CL, Moran AI, Vinciguerra V. Relationship between level of spinal cord injury

- and posttraumatic stress disorder symptoms. Ann N Y Acad Sci. 1997;821:430-432.
- 8. Boldt I, Eriks-Hoogland I, Brinkhof MW, de Bie R, Joggi D, von Elm E. Non-pharmacological interventions for chronic pain in people with spinal cord injury. Cochrane Database of Systematic Reviews. 2014;11:CD009177.
- 9. Bombardier CH, Adams LM, Fann JR, Hoffman JM. Depression Trajectories During the First Year After Spinal Cord Injury. Arch Phys Med Rehabil. 2016;97(2):196-203.
- 10. Bombardier CH, Fann JR, Tate DG, et al. An exploration of modifiable risk factors for depression after spinal cord injury: which factors should we target? Arch Phys Med Rehabil. 2012;93(5):775-781.
- 11. Bombardier CH, Fann JR, Wilson CS, et al. A randomized controlled trial of venlafaxine XR for major depressive disorder after spinal cord injury: Methods and lessons learned. Journal of Spinal Cord Medicine. 2014;37(3):247-263.
- 12. Bombardier CH, Kalpakjian CZ, Graves DE, Dyer JR, Tate DG, Fann JR. Validity of the Patient Health Questionnaire-9 in assessing major depressive disorder during inpatient spinal cord injury rehabilitation. Arch Phys Med Rehabil. 2012;93(10):1838-1845.
- 13. Bombardier CH, Richards JS, Krause JS, Tulsky D, Tate DG. Symptoms of major depression in people with spinal cord injury: implications for screening. Arch Phys Med Rehabil. 2004;85(11):1749-1756.
- 14. Bradley MB. The effect of participating in a functional electrical stimulation exercise program on affect in people with spinal cord injuries. Arch Phys Med Rehabil. 1994;75(6):676-679.
- 15. Burns SM, Boyd BL, Hill J, Hough S. Psychosocial predictors of employment status among men living with spinal cord injury. Rehabilitation Psychology. 2010;55 (1):81-90.
- 16. Burns SM, Hough S, Boyd BL, Hill J. Sexual desire and depression following spinal cord injury: Masculine sexual prowess as a moderator. Sex Roles. 2009;61(1-2):120-129.
- 17. Cao Y, DiPiro ND, Xi J, Krause JS. Unmet expectations of adjustment and depressive symptoms

- among people with chronic traumatic spinal cord injury. Rehabilitation Psychology. 2014;59(3):313-320.
- 18. Cao Y, Massaro JF, Krause JS, Chen Y, Devivo MJ. Suicide mortality after spinal cord injury in the United States: injury cohorts analysis. Arch Phys Med Rehabil. 2014;95(2):230-235.
- 19. Cardenas DD, Nieshoff EC, Suda K, et al. A randomized trial of pregabalin in patients with neuropathic pain due to spinal cord injury. Neurology. 2013;80 (6):533-539.
- 20. Cardenas DD, Warms CA, Turner JA, Marshall H, Brooke MM, Loeser JD. Efficacy of amitriptyline for relief of pain in spinal cord injury: Results of a randomized controlled trial. Pain. 2002;96(3):365-373.
- 21. Catalano D, Chan F, Wilson L, Chiu CY, Muller VR. The buffering effect of resilience on depression among individuals with spinal cord injury: a structural equation model. Rehabilitation Psychology. 2011;56 (3):200-211.
- 22. Charlifue SW, Gerhart KA. Behavioral and demographic predictors of suicide after traumatic spinal cord injury. Arch Phys Med Rehabil. 1991;72(7):488-492.
- 23. Cragg JJ, Noonan VK, Noreau L, Borisoff JF, Kramer JK. Neuropathic pain, depression, and cardio-vascular disease: a national multicenter study. Neuroepidemiology. 2015;44(3):130-137.
- 24. Craig A, Nicholson Perry K, Guest R, Tran Y, Middleton J. Adjustment following chronic spinal cord injury: Determining factors that contribute to social participation. Br J Health Psychol. 2015;20(4):807-823.
- 25. Cuff L, Fann JR, Bombardier CH, Graves DE, Kalpakjian CZ. Depression, pain intensity, and interference in acute spinal cord injury. Topics in Spinal Cord Injury Rehabilitation. 2014;20(1):32-39.
- 26. Danner G, Radnitz CL. Protective factors and post-traumatic stress disorder in veterans with spinal cord injury. International Journal of Rehabilitation & Health. 2000;5(3):195-203.
- 27. Daverat P, Petit H, Kemoun G, Dartigues JF, Barat M. The long term outcome in 149 patients with spinal cord injury. Paraplegia. 1995;33(11):665-668.

- 28. DeVivo MJ, Black KJ, Richards JS, Stover SL. Suicide fol- lowing spinal cord injury. Paraplegia. 1991;29(9):620-627.
- 29. Diego MA, Field T, Hernandez-Reif M, et al. Spinal cord patients benefit from massage therapy. The International journal of neuroscience. 2002;112(2):133-142.
- 30. Dodd Z, Driver S, Warren AM, Riggs S, Clark M. Effects of Adult Romantic Attachment and Social Support on Resilience and Depression in Individuals with Spinal Cord Injuries. Topics in Spinal Cord Injury Rehabilitation. 2015;21(2):156-165.
- 31. Dorstyn D, Mathias J, Denson L. Efficacy of cognitive behavior therapy for the management of psychological outcomes following spinal cord injury: a meta-analysis. J Health Psychol. 2011;16(2):374-391.
- 32. Dorstyn D, Mathias J, Denson L. Applications of telecounselling in spinal cord injury rehabilitation: a systematic review with effect sizes. Clinical Rehabilitation. 2013;27(12):1072-1083.
- 33. Dryden DM, Saunders L, Rowe BH, et al. Depression following Traumatic Spinal Cord Injury. Neuroepidemiology. 2005;25(2):55-61.
- 34. Fann JR, Bombardier CH, Richards J, et al. Venlafaxine extended-release for depression following spinal cord injury: A randomized clinical trial. JAMA Psychiatry. 2015;72(3):247-258.
- 35. Fann JR, Crane DA, Graves DE, Kalpakjian CZ, Tate DG, Bombardier CH. Depression treatment preferences after acute traumatic spinal cord injury. Arch Phys Med Rehabil. 2013;94(12):2389-2395.
- 36. Geisler WO, Jousse AT, Wynne-Jones M, Breithaupt D. Survival in traumatic spinal cord injury. Paraplegia. 1983;21(6):364-373.
- 37. Geyh S, Nick E, Stirnimann D, et al. Self-efficacy and self-esteem as predictors of participation in spinal cord in- jury-an ICF-based study. Spinal Cord. 2012;50(9):699-706.
- 38. Ginis KA, Latimer AE, McKechnie K, et al. Using exercise to enhance subjective well-being among people with spinal cord injury: The mediating influences of stress and pain. Rehabilitation Psychology. 2003;48 (3):157-164.

- 39. Gioia MC, Cerasa A, Di Lucente L, Brunelli S, Castellano V, Traballesi M. Psychological impact of sports activity in spinal cord injury patients. Scand J Med Sci Sports. 2006;16(6):412-416.
- 40. Hagen EM, Lie SA, Rekand T, Gilhus NE, Gronning M. Mortality after traumatic spinal cord injury: 50 years of follow-up. J Neurol Neurosurg Psychiatry. 2010;81(4):368-373.
- 41. Hartkopp A, Bronnum-Hansen H, Seidenschnur AM, Biering-Sorensen F. Survival and cause of death after traumatic spinal cord injury. A long-term epidemiological survey from Denmark.[Erratum appears in Spinal Cord 197 Dec;35(12):862-4]. Spinal Cord. 1997;35(2):76-85.
- 42. Hartoonian N, Hoffman JM, Kalpakjian CZ, Taylor HB, Krause JK, Bombardier CH. Evaluating a spinal cord injury-specific model of depression and quality of life. Arch Phys Med Rehabil. 2014;95 (3):455-465.
- 43. Hassanpour K, Hotz-Boendermaker S, Dokladal P, European Multicenter Study for Human Spinal Cord Injury Study g, Curt A. Low depressive symptoms in acute spinal cord injury compared to other neurological disorders. J Neurol. 2012;259(6):1142-1150.
- 44. Hatcher MB, Whitaker C, Karl A. What predicts post-traumatic stress following spinal cord injury? Br J Health Psychol. 2009;14(Pt 3):541-561.
- 45. Hicks AL, Martin KA, Ditor DS, et al. Long-term exercise training in persons with spinal cord injury: effects on strength, arm ergometry performance and psychological well-being. Spinal cord. 2003;41(1):34-43.
- 46. Hoffman JM, Bombardier CH, Graves DE, Kalpakjian CZ, Krause JS. A longitudinal study of depression from 1 to 5 years after spinal cord injury. Arch Phys Med Rehabil. 2011;92(3):411-418.
- 47. Holicky R, Charlifue S. Ageing with spinal cord injury: the impact of spousal support. Disabil Rehabil. 1999;21(5-6):250-257.
- 48. Houlihan BV, Jette A, Friedman RH, et al. A pilot study of a telehealth intervention for persons with spinal cord dysfunction. Spinal cord. 2013;51(9):715-720.

- 49. Houlihan BV, Jette A, Paasche-Orlow M, et al. A tel-erehabilitation intervention for persons with spinal cord dysfunction. Am J Phys Med Rehabil. 2011;90(9):756-764.
- 50. Hwang M, Chlan KM, Vogel LC, Zebracki K. Substance use in young adults with pediatric-onset spinal cord injury. Spinal Cord. 2012;50(7):497-501.
- 51. Hwang M, Zebracki K, Chlan KM, Vogel LC. Longitudinal employment outcomes in adults with pediatric-onset spinal cord injury. Spinal Cord. 2014;52(6):477-482.
- 52. January AM, Zebracki K, Chlan KM, Vogel LC. Men- tal health and risk of secondary medical complications in adults with pediatric-onset spinal cord injury. Topics in Spinal Cord Injury Rehabilitation. 2014;20 (1):1-12.
- 53. January AM, Zebracki K, Chlan KM, Vogel LC. Symptoms of depression over time in adults with pediatric-onset spinal cord injury. Arch Phys Med Rehabil. 2014;95(3):447-454.
- 54. January AM, Zebracki K, Chlan KM, Vogel LC. Sleep, well-being, and psychological symptoms in adults with pediatric-onset spinal cord injury. Rehabilitation Psychology. 2015;60(4):328-334.
- 55. January AM, Zebracki K, Czworniak A, Chlan KM, Vogel LC. Predictive factors of hospitalization in adults with pediatric-onset SCI: a longitudinal analysis. Spinal Cord. 2015;53(4):314-319.
- 56. Jensen MP, Smith AE, Bombardier CH, Yorkston KM, Miro J, Molton IR. Social support, depression, and physical disability: age and diagnostic group effects. Disabil Health J. 2014;7(2):164-172.
- 57. Jones K, Simpson GK, Briggs L, Dorsett P. Does spirituality facilitate adjustment and resilience among individuals and families after SCI? Disabil Rehabil. 2016;38(10):921-935.
- 58. Kalpakjian CZ, Albright KJ. An examination of depression through the lens of spinal cord injury. Comparative prevalence rates and severity in women and men. Womens Health Issues. 2006;16(6):380-388.
- 59. Kalpakjian CZ, Bombardier CH, Schomer K, Brown PA, Johnson KL. Measuring depression in

- persons with spinal cord injury: a systematic review. Journal of Spinal Cord Medicine. 2009;32(1):6-24.
- 60. Kalpakjian CZ, Houlihan B, Meade MA, et al. Marital status, marital transitions, well-being, and spinal cord injury: an examination of the effects of sex and time. Arch Phys Med Rehabil. 2011;92(3):433-440.
- 61. Kemp B, Krause J, Adkins R. Depression among African Americans, Latinos, and Caucasians with spinal cord injury: A exploratory study. Rehabilitation Psychology. 1999;44(3):235-247.
- 62. Kemp B, Tsukerman D, Kahan J, Adkins R. Predict- ing psychosocial outcomes using a brief measure of quality of life in a sample of people with spinal cord injury. Topics in Spinal Cord Injury Rehabilitation. 2014;20(3):191-196.
- 63. Kennedy P, Taylor N, Hindson L. A pilot investigation of a psychosocial activity course for people with spinal cord injuries. Psychol Health Med. 2006;11(1):91-99.
- 64. Krause J, Coker J, Charlifue S, Whiteneck GG. Depression and subjective well-being among 97 American Indians with spinal cord injury: A descriptive study. Rehabilitation Psychology. 1999;44(4):354-372.
- 65. Krause JS, Broderick L. Outcomes after spinal cord injury: comparisons as a function of gender and race and ethnicity. Arch Phys Med Rehabil. 2004;85 (3):355-362.
- 66. Krause JS, Carter RE, Pickelsimer EE, Wilson D. A prospective study of health and risk of mortality after spinal cord injury. Arch Phys Med Rehabil. 2008;89 (8):1482-1491.
- 67. Krause JS, Coker JL, Charlifue S, Whiteneck GG. Health outcomes among American Indians with spinal cord injury. Arch Phys Med Rehabil. 2000;81 (7):924-931.
- 68. Krause JS, Kemp B, Coker J. Depression after spinal cord injury: relation to gender, ethnicity, aging, and socioeconomic indicators. Arch Phys Med Rehabil. 2000;81(8):1099-1109.
- 69. Krause JS, Saunders LL. Health, secondary conditions, and life expectancy after spinal cord injury. Arch Phys Med Rehabil. 2011;92(11):1770-1775.

- 70. Krause JS, Saunders LL, Newman S. Posttraumatic stress disorder and spinal cord injury. Arch Phys Med Rehabil. 2010;91(8):1182-1187.
- 71. Krause JS, Saunders LL, Reed KS, Coker J, Zhai Y, Johnson E. Comparison of the Patient Health Questionnaire and the Older Adult Health and Mood Questionnaire for self-reported depressive symptoms after spinal cord injury. Rehabilitation Psychology. 2009:54(4):440-448.
- 72. LaVela SL, Burns SP, Goldstein B, Miskevics S, Smith B, Weaver FM. Dysfunctional sleep in persons with spinal cord injuries and disorders. Spinal Cord. 2012;50(9):682-685.
- 73. Le J, Dorstyn D. Anxiety prevalence following spinal cord injury: a meta-analysis. Spinal Cord. 2016;54(8):570-578.
- 74. Lidal IB, Jensen AE, Larsen TW, Stanghelle JK. Fatigue in persons who have lived with spinal cord injury for >20 years. Spinal Cord. 2013;51(2):103-108.
- 75. Lidal IB, Snekkevik H, Aamodt G, Hjeltnes N, Biering- Sorensen F, Stanghelle JK. Mortality after spinal cord injury in Norway. Journal of Rehabilitation Medicine. 2007;39(2):145-151.
- 76. Lin MR, Hwang HF, Yu WY, Chen CY. A prospective study of factors influencing return to work after traumatic spinal cord injury in Taiwan. Arch Phys Med Rehabil. 2009;90(10):1716-1722.
- 77. Liu CW, Huang CC, Chen CH, Yang YH, Chen TW, Huang MH. Prediction of severe neurogenic bowel dysfunction in persons with spinal cord injury. Spinal Cord. 2010;48(7):554-559.
- 78. Lundqvist C, Siosteen A, Sullivan L, Blomstrand C, Lind B, Sullivan M. Spinal cord injuries: a shortened measure of function and mood. Spinal Cord. 1997;35 (1):17-21.
- 79. Lusilla-Palacios P, Castellano-Tejedor C. Spinal cord injury and substance use: a systematic review. Adicciones. 2015;27(4):294-310.
- 80. Martin Ginis KA, Latimer AE, McKechnie K, et al. Using exercise to enhance subjective well-being among people with spinal cord injury: The mediating

- influences of stress and pain. Rehabilitation psychology. 2003;48(3):157-164.
- 81. Martz E. Associations and Predictors of Posttraumatic Stress Levels According to Person-Related, Disability-Related, and Trauma-Related Variables Among Individuals With Spinal Cord Injuries. Rehabilitation Psychology. 2005;50(2):149-157.
- 82. McCullumsmith CB, Kalpakjian CZ, Richards JS, et al. Novel risk factors associated with current suicidal ideation and lifetime suicide attempts in individuals with spinal cord injury. Arch Phys Med Rehabil. 2015;96(5):799-808.
- 83. Mehta S, Orenczuk S, Hansen KT, et al. An evidence-based review of the effectiveness of cognitive behavioral therapy for psychosocial issues post-spinal cord injury. Rehabilitation Psychology. 2011;56(1):15-25.
- 84. Mercier HW, Ni P, Houlihan BV, Jette AM. Differential Impact and Use of a Telehealth Intervention by Persons with MS or SCI. Am J Phys Med Rehabil. 2015;94(11):987-999.
- 85. Migliorini C, Tonge B, Taleporos G. Spinal cord injury and mental health. Aust N Z J Psychiatry. 2008;42(4):309-314.
- 86. Migliorini CE, New PW, Tonge BJ. Comparison of depression, anxiety and stress in persons with traumatic and non-traumatic post-acute spinal cord injury. Spinal Cord. 2009;47(11):783-788.
- 87. Monin JK, Schulz R, Martire LM, Connelly D, Czaja SJ. The personal importance of being independent: Associations with changes in disability and depressive symptoms. Rehabilitation Psychology. 2014;59(1):35-41.
- 88. Muller R, Peter C, Cieza A, Geyh S. The role of social support and social skills in people with spinal cord injury--a systematic review of the literature. Spinal Cord. 2012;50(2):94-106.
- 89. Muller R, Peter C, Cieza A, et al. Social skills: a resource for more social support, lower depression levels, higher quality of life, and participation in individuals with spinal cord injury? Arch Phys Med Rehabil. 2015;96(3):447-455.

- 90. Nam HS, Kim HR, Ha TH, Shin HI. Suicidal ideation in Korean persons with spinal cord injury. Spinal Cord. 2013;51(10):789-793.
- 91. Nelson AL, Groer S, Palacios P, et al. Wheelchairrelated falls in veterans with spinal cord injury residing in the community: a prospective cohort study. Arch Phys Med Rehabil. 2010;91(8):1166-1173.
- 92. Nielsen MS. Prevalence of Posttraumatic Stress Disorder in Persons With Spinal Cord Injuries: The Mediating Effect of Social Support. Rehabilitation Psychology. 2003;48(4):289-295.
- 93. Nooijen CF, Stam HJ, Schoenmakers I, et al. Working mechanisms of a behavioural intervention promoting physical activity in persons with subacute spinal cord injury. Journal of Rehabilitation Medicine. 2016;48(7):583-588.
- 94. Norrbrink C, Lundeberg T. Tramadol in neuropathic pain after spinal cord injury: a randomized, double-blind, placebo-controlled trial. Clin J Pain. 2009;25(3):177-184.
- 95. Oh SJ, Shin HI, Paik NJ, Yoo T, Ku JH. Depressive symptoms of patients using clean intermittent catheterization for neurogenic bladder secondary to spinal cord injury. Spinal Cord. 2006;44(12):757-762.
- 96. O'Shea A, Smedema SM. Understanding depressive symptoms among individuals with spinal cord injuries: biopsychosocial perspective. Rehabilitation Counseling Bulletin. 2014;58(1):20-28.
- 97. Pakpour AH, Kumar S, Scheerman JF, Lin CY, Fridlund B, Jansson H. Oral health-related quality of life in Iranian patients with spinal cord injury: A casecontrol study. Injury. 2016;47(6):1345-1352.
- 98. Perkes SJ, Bowman J, Penkala S. Psychological therapies for the management of co-morbid depression following a spinal cord injury: a systematic review. J Health Psychol. 2014;19(12):1597-1612.
- 99. Peter C, Muller R, Post MW, et al. Depression in spinal cord injury: assessing the role of psychological resources. Rehabilitation Psychology. 2015;60(1):67-80.
- 100. Radnitz CL, Hsu L, Tirch DD, et al. A comparison of posttraumatic stress disorder in veterans with

- and without spinal cord injury. Journal of abnormal psychology. 1998;107(4):676-680.
- 101. Radnitz CL, Hsu L, Willard J, et al. Posttraumatic stress disorder in veterans with spinal cord injury: trauma-related risk factors. J Trauma Stress. 1998;11 (3):505-520.
- 102. Riggins MS, Kankipati P, Oyster ML, Cooper RA, Boninger ML. The relationship between quality of life and change in mobility 1 year postinjury in individuals with spinal cord injury. Arch Phys Med Rehabil. 2011;92(7):1027-1033.
- 103. Rintala DH, Holmes SA, Courtade D, Fiess RN, Tastard LV, Loubser PG. Comparison of the effectiveness of amitriptyline and gabapentin on chronic neuropathic pain in persons with spinal cord injury. Archives of physi- cal medicine and rehabilitation. 2007;88 (12):1547-1560.
- 104. Rintala DH, Robinson-Whelen S, Matamoros R. Subjective stress in male veterans with spinal cord injury. J Rehabil Res Dev. 2005;42(3):291-304.
- 105. Ruhe HG, Dekker JJ, Peen J, Holman R, de Jonghe F. Clinical use of the Hamilton Depression Rating Scale: is in- creased efficiency possible? A post hoc comparison of Ham- ilton Depression Rating Scale, Maier and Bech subscales, Clinical Global Impression, and Symptom Checklist-90 scores. Comprehensive psychiatry. 2005;46 (6):417-427.
- 106. Sakakibara BM, Miller WC, Orenczuk SG, Wolfe DL, Team SR. A systematic review of depression and anxiety measures used with individuals with spinal cord injury. Spinal Cord. 2009;47(12):841-851.
- 107. Saunders LL, Dipiro ND, Krause JS, Brotherton S, Kraft S. Risk of Fall-Related Injuries among Ambulatory Participants with Spinal Cord Injury. Topics in Spinal Cord Injury Rehabilitation. 2013;19 (4):259-266.
- 108. Saunders LL, Krause JS. Psychological factors affecting alcohol use after spinal cord injury. Spinal Cord. 2011;49(5):637-642.
- 109. Saunders LL, Krause JS. Behavioral factors related to fatigue among persons with spinal cord injury. Arch Phys Med Rehabil. 2012;93(2):313-318.

- 110. Saunders LL, Krause JS, Focht KL. A longitudinal study of depression in survivors of spinal cord injury. Spinal Cord. 2012;50(1):72-77.
- 111. Schandler SL, Cohen MJ, Vulpe M, Frank SE. Incidence and characteristics of spinal cord injured patients with a family history of alcoholism. Journal of Studies on Alcohol. 1995;56(5):522-527.
- 112. Schonenberg M, Reimitz M, Jusyte A, Maier D, Badke A, Hautzinger M. Depression, posttraumatic stress, and risk factors following spinal cord injury. Int J Behav Med. 2014;21(1):169-176.
- 113. Schulz R, Czaja SJ, Lustig A, Zdaniuk B, Martire LM, Perdomo D. Improving the quality of life of caregivers of persons with spinal cord injury: a randomized controlled trial. Rehabilitation psychology. 2009;54 (1):1-15.
- 114. Siddall PJ, Cousins MJ, Otte A, Griesing T, Chambers R, Murphy TK. Pregabalin in central neuropathic pain associated with spinal cord injury: a placebocontrolled trial. Neurology. 2006;67(10):1792-1800.
- 115. Sweeney TT, Foote JE. Treatment of drug and alcohol abuse in spinal cord injury veterans. Int J Addict. 1982;17(5):897-904.
- 116. Tasiemski T, Brewer BW. Athletic identity, sport participation, and psychological adjustment in people with spinal cord injury. Adapt Phys Act Q. 2011;28 (3):233-250.
- 117. Tate D, Forchheimer M, Maynard F, Dijkers M. Predicting depression and psychological distress in persons with spinal cord injury based on indicators of handicap. Am J Phys Med Rehabil. 1994;73(3):175-183.
- 118. Tate DG, Forchheimer MB, Krause JS, Meade MA, Bombardier CH. Patterns of alcohol and substance use and abuse in persons with spinal cord injury: risk factors and correlates. Arch Phys Med Rehabil. 2004;85(11):1837-1847.
- 119. Tsai IH, Graves DE, Lai CH, Hwang LY, Pompeii LA. Association of internet use and depression among

- the spinal cord injury population. Arch Phys Med Rehabil. 2014;95(2):236-243.
- 120. Ullrich PM, Lincoln RK, Tackett MJ, Miskevics S, Smith BM, Weaver FM. Pain, depression, and health care utilization over time after spinal cord injury. Rehabilitation Psychology. 2013;58(2):158-165.
- 121. Ullrich PM, Smith BM, Blow FC, Valenstein M, Weaver FM. Depression, healthcare utilization, and comorbid psychiatric disorders after spinal cord injury. Journal of Spinal Cord Medicine. 2014;37 (1):40-45.
- 122. van Gorp S, Kessels AG, Joosten EA, van Kleef M, Pati- jn J. Pain prevalence and its determinants after spinal cord injury: a systematic review. Eur J Pain. 2015;19(1):5-14.
- 123. van Leeuwen CM, Kraaijeveld S, Lindeman E, Post MW. Associations between psychological factors and quality of life ratings in persons with spinal cord injury: a systematic review. Spinal Cord. 2012;50 (3):174-187.
- 124. Williams R, Murray A. Prevalence of depression after spinal cord injury: a meta-analysis. Arch Phys Med Rehabil. 2015;96(1):133-140.
- 125. Williams RT, Heinemann AW, Neumann HD, et al. Eval- uating the Psychometric Properties and Responsiveness to Change of 3 Depression Measures in a Sample of Persons With Traumatic Spinal Cord Injury and Major Depressive Disorder. Arch Phys Med Rehabil. 2016;97(6):929-937.
- 126. Williams RT, Wilson CS, Heinemann AW, Lazowski LE, Fann JR, Bombardier CH. Identifying depression severity risk factors in persons with traumatic spinal cord injury. Rehabilitation psychology. 2014;59(1):50-56.
- 127. Yilmaz B, Yasar E, Koroglu Omac O, Goktepe AS, Tan AK. Gabapentin vs. pregabalin for the treatment of neuropathic pain in patients with spinal cord injury: A crossover study. Turkiye Fiziksel Tip ve Rehabilitasyon Dergisi. 2015;61(1):1-5.

Appendix C. Excluded Studies

The following full-text publications were considered for inclusion but failed to meet the criteria for this report.

Exclusion codes

(first reviewer/second reviewer after consensus):

- 2 = Population not included;
- 3 = Intervention not included;
- 4 = Outcome not included;
- 5 = Comparator not included;
- 6 = Study design not included;

- 7 = Not a study (letter, editorial, non-systematic/narrative review);
- 8 = Non-English language;
- 9 = Inadequate duration;
- 10 = Sample size too small for applicable Key Questions;
- 11 = Systematic Review or meta-analysis used as a source document to identify individual studies; 12 = No original data, duplicate data;
- 13 = Individual study in an included systematic review

Citation Number	Citation	Exclusion Code
1	Abeyasinghe NL, de Zoysa P, Bandara KM, Bartholameuz NA, Bandara JM. The prevalence of symptoms of Post-Traumatic Stress Disorder among soldiers with amputation of a limb or spinal injury: a report from a rehabilitation centre in Sri Lanka. Psychol Health Med. 2012;17(3):376-381.	6/6
2	Abramson AS. Modern concepts of management of the patient with spinal cord injury. Arch Phys Med Rehabil. 1967;48(3):113-121.	7/7
3	Abramson CE, McBride KE, Konnyu KJ, Elliott SL, Team SR. Sexual health outcome measures for individuals with a spinal cord injury: a systematic review. Spinal Cord. 2008;46(5):320-324.	4/4
4	Agar E, Kennedy P, King NS. The Role of Negative Cognitive Appraisals in PTSD Symptoms Following Spinal Cord Injuries. Behavioural and Cognitive Psychotherapy. 2006;34(4):437-452.	13/13
5	Akman RY, Coskun Celik E, Karatas M. Sexuality and sexual dysfunction in spinal cord-injured men in Turkey. Turkish Journal of Medical Sciences. 2015;45(4):758-761.	4/4
6	Allen-Wright S. Expanding health promotion to individuals with spinal cord injuries. SCI Nurs. 1999;16(3):85-88.	6/6
7	Allison DJ, Ditor DS. Targeting inflammation to influence mood following spinal cord injury: A randomized clinical trial. Journal of neuroinflammation. 2015;12(1).	3/3
8	Al-Owesie RM, Moussa NM, Robert AA. Anxiety and depression among traumatic spinal cord injured patients. Neurosciences. 2012;17(2):145-150.	4/4
9	Alschuler KN, Kratz AL, Ehde DM. Resilience and vulnerability in individuals with chronic pain and physical disability. Rehabilitation Psychology. 2016;61(1):7-18.	10/10
10	Anderson CJ, Krajci KA, Vogel LC. Life satisfaction in adults with pediatric-onset spinal cord injuries. Journal of Spinal Cord Medicine. 2002;25(3):184-190.	4/4
11	Anderson CJ, Vogel LC, Betz RR, Willis KM. Overview of adult outcomes in pediatric-onset spinal cord injuries: implications for transition to adulthood. Journal of Spinal Cord Medicine. 2004;27 Suppl 1:S98-106.	10/10
12	Anderson CJ, Vogel LC, Chlan KM, Betz RR, McDonald CM. Depression in adults who sustained spinal cord injuries as children or adolescents. Journal of Spinal Cord Medicine. 2007;30 Suppl 1:S76-82.	6/6
13	Anderson CJ, Vogel LC, Chlan KM, Betz RR. Coping with spinal cord injury: strategies used by adults who sustained their injuries as children or adolescents. Journal of Spinal Cord Medicine. 2008;31(3):290-296.	4/4
14	Anton HA, Miller WC, Townson AF. Measuring fatigue in persons with spinal cord injury. Arch Phys Med Rehabil. 2008;89(3):538-542.	10/10
15	Ataoglu E, Tiftik T, Kara M, Tunc H, Ersoz M, Akkus S. Effects of chronic pain on quality of life and depression in patients with spinal cord injury. Spinal Cord. 2013;51(1):23-26.	2/2
16	Avluk OC, Gurcay E, Gurcay AG, Karaahmet OZ, Tamkan U, Cakci A. Effects of chronic pain on function, depression, and sleep among patients with traumatic spinal cord injury. Ann Saudi Med. 2014;34(3):211-216.	10/10
17	Baker FA, Rickard N, Tamplin J, Roddy C. Flow and Meaningfulness as Mechanisms of Change in Self-Concept and Well-Being Following a Songwriting Intervention for People in the Early Phase of Neurorehabilitation. Front Hum Neurosci. 2015;9:299.	3/3
18	Barbonetti A, Cavallo F, Felzani G, Francavilla S, Francavilla F. Erectile dysfunction is the main determinant of psychological distress in men with spinal cord injury. J Sex Med. 2012;9(3):830-836.	10/10
19	Beedie A, Kennedy P. Quality of social support predicts hopelessness and depression post spinal cord injury. Journal of Clinical Psychology in Medical Settings. 2002;9(3):227-234.	13/13
20	Benony H, Daloz L, Bungener C, Chahraoui K, Frenay C, Auvin J. Emotional factors and subjective quality of life in subjects with spinal cord injuries. Am J Phys Med Rehabil. 2002;81(6):437-445.	6/6
21	Bermond B, Schuerman J, Nieuwenhuijse B, Fasotti L, et al. Spinal cord lesions: Coping and mood states. Clinical Rehabilitation. 1987;1(2):111-117.	10/10

Continued

Citation Number	Citation	Exclusion Code
22	Berry JW, Elliott TR, Rivera P. Resilient, undercontrolled, and overcontrolled personality prototypes among persons with spinal cord injury. J Pers Assess. 2007;89(3):292-302.	13/13
23 24	Biering-Sorensen F, Hartkopp A. Spinal cord lesions. Curr Opin Neurol. 1995;8(6):451-455. Blanes L, Carmagnani MI, Ferreira LM. Quality of life and self-esteem of persons with paraplegia	6/6 10/10
25	living in Sao Paulo, Brazil. Qual Life Res. 2009;18(1):15-21. Boakye M, Moore R, Kong M, Skirboll SL, Arrigo RT. Health-related quality-of-life status in Veterans with spinal disorders. Quality of Life Research: An International Journal of Quality of Life Aspects of	2/2
26	Treatment, Care & Rehabilitation. 2013;22(1):45-52. Bockian NR, Lee A, Fidanque CS. Personality disorders and spinal cord injury: A pilot study. Journal of Clinical Psychology in Medical Settings. 2003;10(4):307-313.	4/4
27	Boekamp JR, Overholser JC, Schubert DS. Depression following a spinal cord injury. Int J Psychiatry Med. 1996;26(3):329-349.	11/11
28	Bombardier CH, Ehde DM, Stoelb B, Molton IR. The relationship of age-related factors to psychological functioning among people with disabilities. Phys Med Rehabil Clin N Am. 2010;21(2):281-297.	11/11
29	Bombardier CH, Rimmele CT. Alcohol use and readiness to change after spinal cord injury. Arch Phys Med Rehabil. 1998;79(9):1110-1115.	10/10
30	Bombardier CH, Stroud MW, Esselman PC, Rimmele CT. Do preinjury alcohol problems predict poorer rehabilitation progress in persons with spinal cord injury? Arch Phys Med Rehabil. 2004;85(9):1488-1492.	13/13
31 32	Bombardier CH. Depression and spinal cord injury. Arch Phys Med Rehabil. 2014;95(2):413-414. Bombardier CH. Venlafaxine XR for depression and pain in spinal cord injury: Major outcomes. Journal of spinal cord medicine.36(5 SPEC. ISSUE):551.	7/7 7/7
33	Bonanno GA, Kennedy P, Galatzer-Levy IR, Lude P, Elfstrom ML. Trajectories of resilience, depression, and anxiety following spinal cord injury. Rehabilitation Psychology. 2012;57(3):236-247.	4/4
34	Borsbo B, Gerdle B, Peolsson M. Impact of the interaction between self-efficacy, symptoms and catastrophising on disability, quality of life and health in with chronic pain patients. Disabil Rehabil. 2010;32(17):1387-1396.	10/10
35	Boxall J. Care of the spinal cord-injured patient. Can Fam Physician. 1989;35:2435-2439.	7/7
36	Bozzacco V. Vulnerability and alcohol and substance abuse in spinal cord injury. Rehabil Nurs. 1990;15(2):70-72.	7/7
37	Bracken MB, Shepard MJ. Coping and adaptation following acute spinal cord injury: a theoretical analysis. Paraplegia. 1980;18(2):74-85.	7/7
38	Buckelew SP, Burk JP, Brownlee-Duffeck M, Frank RG, DeGood D. Cognitive and somatic aspects of depression among a rehabilitation sample: Reliability and validity of SCL-90-R research subscales. Rehabilitation Psychology. 1988;33(2):67-75.	4/4
39	Buckelew SP, Frank RG, Elliott TR, Chaney J, Hewett J. Adjustment to spinal cord injury: stage theory revisited. Paraplegia. 1991;29(2):125-130.	13/13
40	Budh CN, Osteraker AL. Life satisfaction in individuals with a spinal cord injury and pain. Clinical Rehabilitation. 2007;21(1):89-96.	10/10
41	Buunk AP, Zurriaga R, Gonzalez P. Social comparison, coping and depression in people with spinal cord injury. Psychol Health. 2006;21(6):791-807.	13/13
42	Byra S. Posttraumatic growth in people with traumatic long-term spinal cord injury: predictive role of basic hope and coping. Spinal Cord. 2016;54(6):478-482.	4/4
43	Cairns DM, Adkins RH, Scott MD. Pain and depression in acute traumatic spinal cord injury: origins of chronic problematic pain? Arch Phys Med Rehabil. 1996;77(4):329-335.	10/10
44	Cao Y, Walker EA, Krause JS. Environmental barriers and subjective health among people with chronic spinal cord injury: A cohort study. Journal of Spinal Cord Medicine. 2015;38(4):526-531.	2/2
45	Castriotta RJ, Wilde MC, Sahay S. Sleep disorders in spinal cord injury. Sleep Medicine Clinics. 2012;7(4):643-653.	7/7
46	Chan RC, Lee PW, Lieh-Mak F. Coping with spinal cord injury: personal and marital adjustment in the Hong Kong Chinese setting. Spinal Cord. 2000;38(11):687-696.	10/10
47	Chan RC, Lee PW, Lieh-Mak F. The pattern of coping in persons with spinal cord injuries. Disabil Rehabil. 2000;22(11):501-507.	13/13
48	Chapin MH, Holbert D. Differences in affect, life satisfaction, and depression between successfully and unsuccessfully rehabilitated persons with spinal cord injuries. Rehabilitation Counseling Bulletin. 2009;53(1):6-15.	10/10
49	Charlifue S, Apple D, Burns SP, et al. Mechanical ventilation, health, and quality of life following spinal cord injury. Archives of physical medicine and rehabilitation. 2011;92(3):457-463.	4/4
50	Charlifue S, Gerhart K. Changing psychosocial morbidity in people aging with spinal cord injury. Neurorehabilitation. 2004;19(1):15-23.	2/2
51	Chen Y, Anderson CJ, Vogel LC, Chlan KM, Betz RR, McDonald CM. Change in life satisfaction of adults with pediatric-onset spinal cord injury. Arch Phys Med Rehabil. 2008;89(12):2285-2292.	10/10

Citation Number	Citation	Exclusion Code
52	Chevalier Z, Kennedy P, Sherlock O. Spinal cord injury, coping and psychological adjustment: a literature review. Spinal Cord. 2009;47(11):778-782.	7/7
53	Chung MC, Preveza E, Papandreou K, Prevezas N. Locus of control among spinal cord injury patients with different levels of posttraumatic stress disorder. Psychiatry Res. 2007;152(2-3):253-260.	10/10
54	Chung MC, Preveza E, Papandreou K, Prevezas N. Spinal cord injury, posttraumatic stress, and locus of control among the elderly: a comparison with young and middle-aged patients. Psychiatry. 2006;69(1):69-80.	10/10
55	Chung MC, Preveza E, Papandreou K, Prevezas N. The relationship between posttraumatic stress disorder following spinal cord injury and locus of control. J Affect Disord. 2006;93(1-3):229-232.	2/2
56	Clay DL, Hagglund KJ, Frank RG, Elliott TR, Chaney JM. Enhancing the accuracy of depression diagnosis in patients with spinal cord injury using Bayesian analysis. Rehabilitation Psychology. 1995;40(3):171-180.	13/13
57	Cobo Cuenca AI, Sampietro-Crespo A, Virseda-Chamorro M, Martin-Espinosa N. Psychological impact and sexual dysfunction in men with and without spinal cord injury. J Sex Med. 2015;12(2):436-444.	4/4
58	Coleman JA, Harper LA, Perrin PB, et al. The relationship between physical and mental health variables in individuals with spinal cord injury from Latin America. Pm R. 2015;7(1):9-16.	2/2
59	Cook DW. A multivariate analysis of motivational attributes among spinal cord injured rehabilitation clients. International Journal of Rehabilitation Research. 1981;4(1):5-15.	4/4
60	Cook DW. Psychological adjustment to spinal cord injury: Incidence of denial, depression, and anxiety. Rehabilitation Psychology. 1979;26(3):97-104.	4/4
61	Cook DW. Psychological aspects of spinal cord injury. Rehabilitation Counseling Bulletin. 1976;19(4):535-543.	7/7
62	Cook KF, Bamer AM, Amtmann D, Molton IR, Jensen MP. Six patient-reported outcome measurement information system short form measures have negligible age- or diagnosis-related differential item functioning in individuals with disabilities. Arch Phys Med Rehabil. 2012;93(7):1289-1291.	2/2
63	Cook KF, Kallen MA, Bombardier C, et al. Do measures of depressive symptoms function differently in people with spinal cord injury versus primary care patients: The ces-d, phq-9, and promis-d. Quality of Life Research: An International Journal of Quality of Life Aspects of Treatment, Care & Rehabilitation. 2016:No Pagination Specified.	4/4
64	Corcoran JR, Frank RG, Elliott TR. The interpersonal influence of depression following spinal cord injury: A methodological study. Journal of the Multihandicapped Person. 1988;1(3):161-174.	6/6
65	Coyle CP, Lesnik-Emas S, Kinney WB. Predicting life satisfaction among adults with spinal cord injuries. Rehabilitation Psychology. 1994;39(2):95-112.	13/13
66	Coyle CP, Shank JW, Kinney W, Hutchins DA. Psychosocial functioning and changes in leisure lifestyle among individuals with chronic secondary health problems related to spinal cord injury. Therapeutic Recreation Journal. 1993;27(4):239-252.	13/13
67	Craig A, Hancock K, Dickson H. Improving the long-term adjustment of spinal cord injured persons. Spinal cord. 1999;37(5):345-350.	10/10
68	Craig A, Hancock K. Difficulities in implementing group cognitive behaviour therapy for spinal cord injured persons: A clinical discussion. Australian Psychologist. 1994;29(2):98-102.	7/7
69	Craig A, Nicholson Perry K, Guest R, et al. Prospective Study of the Occurrence of Psychological Disorders and Comorbidities after Spinal Cord Injury. Archives of physical medicine and rehabilitation. 2015;96(8):1426-1434.	10/10
70	Craig A, Rodrigues D, Tran Y, Guest R, Bartrop R, Middleton J. Developing an algorithm capable of discriminating depressed mood in people with spinal cord injury. Spinal Cord. 2014;52(5):413-416.	4/4
71	Craig A, Tran Y, Middleton J. Psychological morbidity and spinal cord injury: a systematic review. Spinal Cord. 2009;47(2):108-114.	2/2
72	Craig A, Tran Y, Siddall P, et al. Developing a model of associations between chronic pain, depressive mood, chronic fatigue, and self-efficacy in people with spinal cord injury. Journal of Pain. 2013;14(9):911-920.	4/4
73	Craig A, Tran Y, Wijesuriya N, Middleton J. Fatigue and tiredness in people with spinal cord injury. J Psychosom Res. 2012;73(3):205-210.	10/10
74	Craig AR, Hancock K, Chang E, Dickson H. Immunizing against depression and anxiety after spinal cord injury. Archives of physical medicine and rehabilitation. 1998;79:375-377.	13/13
75	Craig AR, Hancock K, Dickson H, Chang E. Long-term psychological outcomes in spinal cord injured persons: results of a controlled trial using cognitive behavior therapy. Archives of physical medicine and rehabilitation. 1997;78(1):33-38.	13/13
76	Craig AR, Hancock KM, Dickson HG. A longitudinal investigation into anxiety and depression in the first 2 years following a spinal cord injury. Paraplegia. 1994;32(10):675-679.	10/10
77	Craig AR, Hancock KM, Dickson HG. Spinal cord injury: a search for determinants of depression two years after the event. Br J Clin Psychol. 1994;33(Pt 2):221-230.	13/13

Citation Number	Citation	Exclusion Code
78	Craven C, Hitzig SL, Mittmann N. Impact of impairment and secondary health conditions on health preference among Canadians with chronic spinal cord injury. Journal of Spinal Cord Medicine. 2012;35(5):361-370.	10/10
79	Crawford JJ, Gayman AM, Tracey J. An examination of post-traumatic growth in Canadian and American ParaSport athletes with acquired spinal cord injury. Psychology of Sport and Exercise. 2014;15(4):399-406.	10/10
80	Crewe NM. Life stories of people with long-term spinal cord injury. Rehabilitation Counseling Bulletin. 1997;41(1):26-42.	6/6
81	Crews WD, Jr., Hensley LG, Goering AM, Barth JT, Rusek JT. Spinal cord injury and anxiety: a comprehensive review. Neurorehabilitation. 1998;11(3):155-174.	11/11
82	Crisp R. Depression and occupational disability in five diagnostic groups: a review of recent research. Disabil Rehabil. 2007;29(4):267-279.	11/11
83	Crisp R. The long-term adjustment of 60 persons with spinal cord injury. Australian Psychologist. 1992;27(1):43-47.	10/10
84	Crumbaugh JC. Cross-validation of Purpose-in-Life test based on Frankl's concepts. Journal of individual psychology. 1968;24(1):74-81.	6/6
85	Cruz-Almeida Y, Alameda G, Widerstrom-Noga EG. Differentiation between pain-related interference and interference caused by the functional impairments of spinal cord injury. Spinal Cord. 2009;47(5):390-395.	4/4
86	Cushman LA, Dijkers MP. Depressed mood in spinal cord injured patients: staff perceptions and patient realities. Arch Phys Med Rehabil. 1990;71(3):191-196.	4/4
87	Dahlberg A, Alaranta H, Sintonen H. Health-related quality of life in persons with traumatic spinal cord lesion in Helsinki. Journal of Rehabilitation Medicine. 2005;37(5):312-316.	2/2
88	Dajpratham P, Kongkasuwan R. Quality of life among the traumatic spinal cord injured patients. J Med Assoc Thai. 2011;94(10):1252-1259.	10/10
89	Daniel A, Manigandan C. Efficacy of leisure intervention groups and their impact on quality of life among people with spinal cord injury. International journal of rehabilitation research. 2005;Internationale Zeitschrift für Rehabilitationsforschung. Revue internationale de recherches de	4/4
90	réadaptation. 28(1):43-48. Davidoff G, Roth E, Thomas P, et al. Depression among acute spinal cord injury patients: A study utilizing the Zung Self-Rating Depression Scale. Rehabilitation Psychology. 1990;35(3):171-180.	10/10
91	Davidoff G, Roth E, Thomas P, et al. Depression and neuropsychological test performance in acute spinal cord injury patients: lack of correlation. Arch Clin Neuropsychol. 1990;5(1):77-88.	10/10
92	Davidoff GN, Roth EJ, Richards JS. Cognitive deficits in spinal cord injury: epidemiology and outcome. Arch Phys Med Rehabil. 1992;73(3):275-284.	6/6
93	Davis F, Kumaran D S, Nair S, J V. Physical fitness training for people with spinal cord injury. Cochrane Database of Systematic Reviews. 2009(4).	11/11
94	de Carvalho SA, Andrade MJ, Tavares MA, de Freitas JL. Spinal cord injury and psychological response. Gen Hosp Psychiatry. 1998;20(6):353-359.	10/10
95	Dean RE, Kennedy P. Measuring appraisals following acquired spinal cord injury: a preliminary psychometric analysis of the appraisals of disability. Rehabilitation Psychology. 2009;54(2):222-231.	2/2
96	Decker SD, Schulz R. Correlates of life satisfaction and depression in middle-aged and elderly spinal cord-injured persons. Am J Occup Ther. 1985;39(11):740-745.	10/10
97	deRoon-Cassini TA, de S, Valvano AK, Hastings J, Brasel KJ. Meaning-Making Appraisals Relevant to Adjustment for Veterans With Spinal Cord Injury. Psychol Serv. 2012:No Pagination Specified.	10/10
98	Derrett S, Beaver C, Sullivan MJ, Herbison GP, Acland R, Paul C. Traumatic and non-traumatic spinal cord impairment in New Zealand: incidence and characteristics of people admitted to spinal units. In Prev. 2012;18(5):343-346.	10/10
99	DeVivo MJ, Kartus PL, Stover SL, Rutt RD, Fine PR. Cause of death for patients with spinal cord injuries. Arch Intern Med. 1989;149(8):1761-1766.	4/4
100	Dijkers M, Cushman LA. Differences between rehabilitation disciplines in views of depression in spinal cord injury patients. Paraplegia. 1990;28(6):380-391.	3/3
101	Dijkers MP. Quality of life of individuals with spinal cord injury: a review of conceptualization, measurement, and research findings. J Rehabil Res Dev. 2005;42(3 Suppl 1):87-110.	7/7
102	Dinoff BL, Richards JS, Ness TJ. Use of topiramate for spinal cord injury-related pain. Journal of Spinal Cord Medicine. 2003;26(4):401-403.	6/6
103	Dishman RK, Berthoud HR, Booth FW, et al. Neurobiology of exercise. Obesity (Silver Spring). 2006;14(3):345-356.	6/6
104	Ditre JW, Radnitz CL. Pre- and Postinjury Substance Misuse Among Veterans With Spinal Cord Injury. Rehabilitation Psychology. 2005;50(2):142-148.	10/10
105	Dorstyn D, Mathias J, Denson L, Robertson M. Effectiveness of telephone counseling in managing psychological outcomes after spinal cord injury: a preliminary study. Archives of physical medicine and rehabilitation. 2012;93(11):2100-2108.	13/13

Continued

Citation Number	Citation	Exclusion Code
106	Dorstyn D, Mathias J, Denson L. Psychosocial outcomes of telephone-based counseling for adults with an acquired physical disability: A meta-analysis. Rehabilitation Psychology. 2011;56(1):1-14.	11/11
107	Drainoni ML, Houlihan B, Williams S, et al. Patterns of Internet use by persons with spinal cord injuries and relationship to health-related quality of life. Arch Phys Med Rehabil. 2004;85(11):1872-1879.	4/4
108	Driver S, Warren AM, Reynolds M, et al. Identifying predictors of resilience at inpatient and 3-month post-spinal cord injury. Journal of Spinal Cord Medicine. 2016;39(1):77-84.	10/10
109	Ducharme S, Murray FM. The Role of Self Destruction in spinal cord injury mortality. Model Systems SCI Digest. 1980;2:29-38.	10/10
110	Duchnick JJ, Letsch EA, Curtiss G. Coping effectiveness training during acute rehabilitation of spinal cord injury/dysfunction: a randomized clinical trial. Rehabilitation psychology. 2009;54(2):123-132.	13/13
111	Dunn J, Sinnott KA, Nunnerley J, Scheuringer M. Utilisation of patient perspective to validate clinical measures of outcome following spinal cord injury. Disabil Rehabil. 2009;31(12):967-975.	4/4
112	Dunn M, Davis R. The perceived effects of marijuana on spinal cord injured males. Paraplegia. 1974;12(3):175.	6/6
113	Dunn M, Love L, Ravesloot C. Subjective health in spinal cord injury after outpatient healthcare follow-up. Spinal Cord. 2000;38(2):84-91.	2/2
114	Ebrahimzadeh MH, Makhmalbaf H, Soltani-Moghaddas SH, Mazloumi SM. The spinal cord injury quality-of-life-23 questionnaire, Iranian validation study. J. 2014;19(4):349-354.	4/4
115	Ebrahimzadeh MH, Soltani-Moghaddas SH, Birjandinejad A, Omidi-Kashani F, Bozorgnia S. Quality of life among veterans with chronic spinal cord injury and related variables. Arch. 2014;3(2):e17917.	10/10
116	Egner A, Phillips VL, Vora R, Wiggers E. Depression, fatigue, and health-related quality of life among people with advanced multiple sclerosis: results from an exploratory telerehabilitation study. NeuroRehabilitation. 2003;18(2):125-133.	2/2
117	Elliot TR, Kurylo M, Chen Y, Hicken B. Alcohol abuse history and adjustment following spinal cord injury. Rehabilitation Psychology. 2002;47(3):278-290.	2/2
118	Elliott FC. A nursing protocol for anxiety following catastrophic injury. Rehabil Nurs. 1983;8(3):18-20, 38.	6/6
119	Elliott TR, Frank RG. Depression following spinal cord injury. Arch Phys Med Rehabil. 1996;77(8):816-823.	7/7
120	Elliott TR, Godshall FJ, Herrick SM, Witty TE, Spruell M. Problem-solving appraisal and psychological adjustment following spinal cord injury. Cognitive Therapy and Research. 1991;15(5):387-398.	13/13
121	Elliott TR, Herrick SM, Patti AM, Witty TE, Godshall FJ, Spruell M. Assertiveness, social support, and psychological adjustment following spinal cord injury. Behav Res Ther. 1991;29(5):485-493.	13/13
122	Elliott TR, Herrick SM, Witty TE, Godshall F, Spruell M. Social support and depression following spinal cord injury. Rehabilitation Psychology. 1992;37(1):37-48.	13/13
123	Elliott TR, Kennedy P. Treatment of Depression Following Spinal Cord Injury: An Evidence-Based Review. Rehabilitation Psychology. 2004;49(2):134-139.	11/11
124	Elliott TR, Witty TE, Herrick S, Hoffman JT. Negotiating reality after physical loss: hope, depression, and disability. J Pers Soc Psychol. 1991;61(4):608-613. Ernst FA. Contrasting perceptions of distress by research personnel and their spinal cord injured	10/10 4/4
125	subjects. Am J Phys Med. 1987;66(1):12-15.	4/4
26 27	Faber RA. Depression and spinal cord injury. Neuroepidemiology. 2005;25(2):53-54. Fann JR, Bombardier CH, Dikmen S, et al. Validity of the Patient Health Questionnaire-9 in assessing depression following traumatic brain injury. J Head Trauma Rehabil. 2005;20(6):501-511.	7/7 2/2
128	Fann JR, Bombardier CH, Richards JS, et al. Depression after spinal cord injury: comorbidities, mental health service use, and adequacy of treatment. Arch Phys Med Rehabil. 2011;92(3):352-360.	10/10
129	Fedoroff JP, Lipsey JR, Starkstein SE, Forrester A, Price TR, Robinson RG. Phenomenological comparisons of major depression following stroke, myocardial infarction or spinal cord lesions. J Affect Disord. 1991;22(1-2):83-89.	10/10
130	Fekete C, Siegrist J, Reinhardt JD, Brinkhof MW, Swi SCISG. Is financial hardship associated with reduced health in disability? The case of spinal cord injury in Switzerland. PLoS ONE. 2014;9(2):e90130.	2/2
131	Ferington FE. Personal control and coping effectiveness in spinal cord injured persons. Res Nurs Health. 1986;9(3):257-265.	4/4
132	Field-Fote EC, Brown KM, Lindley SD. Influence of posture and stimulus parameters on post- activation depression of the soleus H-reflex in individuals with chronic spinal cord injury. Neurosci Lett. 2006;410(1):37-41.	4/4
133	Findley PA, Banerjea R, Sambamoorthi U. Excess mortality associated with mental illness and substance use disorders among veteran clinic users with spinal cord injury. Disabil Rehabil. 2011;33(17-18):1608-1615.	13/13
134	Fogelberg DJ, Vitiello MV, Hoffman JM, Bamer AM, Amtmann D. Comparison of self- report sleep measures for individuals with multiple sclerosis and spinal cord injury. Arch Phys Med Rehabil. 2015;96(3):478-483.	4/4

Citation Number	Citation	Exclusion Code
135	Foreman PE, Cull J, Kirkby RJ. Sports participation in individuals with spinal cord injury: demographic and psychological correlates. International Journal of Rehabilitation Research. 1997;20(2):159-168.	2/2
136	Frank RG, Chaney JM, Clay DL, et al. Dysphoria: a major symptom factor in persons with disability or chronic illness. Psychiatry Res. 1992;43(3):231-241.	4/4
137	Frank RG, Elliott TR, Buckelew SP, Haut AE. Age as a factor in response to spinal cord injury. Am J Phys Med Rehabil. 1988;67(3):128-131.	10/10
138	Frank RG, Elliott TR, Corcoran JR, Wonderlich SA. Depression after spinal cord injury: Is it necessary? Clinical Psychology Review. 1987;7(6):611-630.	11/11
139	Frank RG, Elliott TR, Wonderlich SA, Corcoran JR, Umlauf RL, Ashkanazi GS. Gender differences in the interpersonal response to depression and spinal cord injury. Cognitive Therapy and Research. 1987;11(4):437-448.	3/3
140	Frank RG, Elliott TR. Life stress and psychologic adjustment following spinal cord injury. Arch Phys Med Rehabil. 1987;68(6):344-347.	10/10
141	Frank RG, Elliott TR. Spinal cord injury and health locus of control beliefs. Paraplegia. 1989;27(4):250-256.	13/13
142	Frank RG, Kashani JH, Wonderlich SA, Lising A, Visot LR. Depression and adrenal function in spinal cord injury. Am J Psychiatry. 1985;142(2):252-253.	10/10
143	Frank RG, Umlauf RL, Wonderlich SA, Askanazi GS, Buckelew SP, Elliott TR. Differences in coping styles among persons with spinal cord injury: A cluster-analytic approach. Journal of Consulting and Clinical Psychology. 1987;55(5):727-731.	10/10
144	Franulic A, Carbonell CG, Pinto P, Sepulveda I. Psychosocial adjustment and employment outcome 2, 5 and 10 years after TBI. Brain Inj. 2004;18(2):119-129.	2/2
145	Fraser AR. The psychiatrist in the spinal unit. Australian and New Zealand Journal of Psychiatry. 1982;16(4):246-250.	7/7
146	Frisbie JH, Tun CG. Drinking and spinal cord injury. J Am Paraplegia Soc. 1984;7(4):71-73.	4/4
147	Fuhrer MJ, Rintala DH, Hart KA, Clearman R, Young ME. Depressive symptomatology in persons with spinal cord injury who reside in the community. Arch Phys Med Rehabil. 1993;74(3):255-260.	4/4
148	Fukunishi I, Koyama I, Tombimatsu H. Psychological acceptance and alexithymia in spinal cord injury patients. Psychol Rep. 1995;76(2):475-481.	4/4
149	Fullerton DT, Harvey RF, Klein MH, Howell T. Psychiatric disorders in patients with spinal cord injuries. Arch Gen Psychiatry. 1981;38(12):1369-1371.	10/10
150	Furlan JC, Bracken MB, Fehlings MG. Is age a key determinant of mortality and neurological outcome after acute traumatic spinal cord injury? Neurobiology of aging. 2010;31(3):434-446.	2/2
151	Galea M, Tumminia J, Garback LM. Telerehabilitation in spinal cord injury persons: a novel approach. Telemed J E Health. 2006;12(2):160-162.	7/7
152	Galvin LR, Godfrey HP. The impact of coping on emotional adjustment to spinal cord injury (SCI): review of the literature and application of a stress appraisal and coping formulation. Spinal Cord. 2001;39(12):615-627.	7/7
153	Gerhart KA, Johnson RL, Whiteneck GG. Health and psychosocial issues of individuals with incomplete and resolving spinal cord injuries. Paraplegia. 1992;30(4):282-287.	13/13
154	Geyh S, Kunz S, Muller R, Peter C, Swi SCISG. Describing functioning and health after spinal cord injury in the light of psychological-personal factors. Journal of Rehabilitation Medicine. 2016;48(2):219-234.	11/11
155	Geyh S, Nick E, Stirnimann D, Ehrat S, Muller R, Michel F. Biopsychosocial outcomes in individuals with and without spinal cord injury: a Swiss comparative study. Spinal Cord. 2012;50(8):614-622.	10/10
156	Giannini MJ, Bergmark B, Kreshover S, Elias E, Plummer C, O'Keefe E. Understanding suicide and disability through three major disabling conditions: Intellectual disability, spinal cord injury, and multiple sclerosis. Disabil Health J. 2010;3(2):74-78.	11/11
157	Giannoccaro MP, Moghadam KK, Pizza F, et al. Sleep disorders in patients with spinal cord injury. Sleep Med Rev. 2013;17(6):399-409.	4/4
158	Giardino ND, Jensen MP, Turner JA, Ehde DM, Cardenas DD. Social environment moderates the association between catastrophizing and pain among persons with a spinal cord injury. Pain. 2003;106(1-2):19-25.	10/10
159	Gichia JW, Glenn L. Sufficiency of a two factor model for posttraumatic stress disorder symptoms in spinal injury. Arch Psychiatr Nurs. 2015;29(4):236-237.	7/7
160	Glass CA. Psychological intervention in physical disability-with special reference to spinal cord injury. Journal of Mental Health. 1994;3(4):467-476.	7/7
161	Goldman RL, Radnitz CL, McGrath RE. Posttraumatic stress disorder and major depression in veterans with spinal cord injury. Rehabilitation Psychology. 2008;53(2):162-170.	2/2
162	Gorman C, Kennedy P, Hamilton LR. Alterations in self-perceptions following childhood onset of spinal cord injury. Spinal Cord. 1998;36(3):181-185.	10/10
163	Graves DE, Bombardier CH. Improving the efficiency of screening for major depression in people with spinal cord injury. Journal of Spinal Cord Medicine. 2008;31(2):177-184.	4/4

Citation Number	Citation	Exclusion Code
164	Griffiths HC, Clinpsy D, Kennedy P. Continuing with life as normal:: positive psychological outcomes following spinal cord injury. Topics in Spinal Cord Injury Rehabilitation. 2012;18(3):241-252.	10/10
165	Griffiths T, Myers D, Talbot A. A study of the validity of the scaled version of the General Health Questionnaire in paralysed spinally injured out-patients. Psychological Medicine. 1993;23(2):497-504.	4/4
166	Groah SL, Stiens SA, Gittler MS, Kirshblum SC, McKinley WO. Spinal cord injury medicine. 5. Preserving wellness and independence of the aging patient with spinal cord injury: a primary care approach for the rehabilitation medicine specialist. Arch Phys Med Rehabil. 2002;83(3 Suppl 1):S82-89, S90-88.	6/6
167	Guest R, Craig A, Perry KN, et al. Resilience following spinal cord injury: A prospective controlled study investigating the influence of the provision of group cognitive behavior therapy during inpatient rehabilitation. Rehabilitation psychology. 2015;60(4):311-321.	6/6
168	Gupta A, Deepika S, Taly AB, Srivastava A, Surender V, Thyloth M. Quality of life and psychological problems in patients undergoing neurological rehabilitation. Ann. 2008;11(4):225-230.	10/10
169	Hagen EM, Grimstad KE, Bovim L, Gronning M. Patients with traumatic spinal cord injuries and their satisfaction with their general practitioner. Spinal Cord. 2012;50(7):527-532.	4/4
170	Hagen EM, Rekand T, Gilhus NE, Gronning M. Traumatic spinal cord injuries incidence, mechanisms and course. Tidsskr Nor Laegeforen. 2012;132(7):831-837.	4/4
171	Hajiaghababaei M, Javidan AN, Saberi H, et al. Female sexual dysfunction in patients with spinal cord injury: a study from Iran. Spinal Cord. 2014;52(8):646-649.	2/2
172	Hammell KW, Miller WC, Forwell SJ, Forman BE, Jacobsen BA. Fatigue and spinal cord injury: a qualitative analysis. Spinal Cord. 2009;47(1):44-49.	4/4
173	Hampton NZ, Qin-Hilliard DB. Dimensions of quality of life for Chinese adults with spinal cord injury: a qualitative study. Disabil Rehabil. 2004;26(4):203-212.	4/4
174	Hancock K, Craig A, Tennant C, Chang E. The influence of spinal cord injury on coping styles and self-perceptions: a controlled study. Aust N Z J Psychiatry. 1993;27(3):450-456.	4/4
75	Hancock KM, Craig AR, Dickson HG, Chang E, Martin J. Anxiety and depression over the first year of spinal cord injury: a longitudinal study. Paraplegia. 1993;31(6):349-357.	10/10
176	Hanley MA, Raichle K, Jensen M, Cardenas DD. Pain catastrophizing and beliefs predict changes in pain interference and psychological functioning in persons with spinal cord injury. The Journal of Pain. 2008;9(9):863-871.	10/10
177	Hanover J. What are the assessment components for SCI patients pertaining to evaluation of substance abuse? SCI Nurs. 1990;7(2):38.	7/7
178	Haran MJ, Lee BB, King MT, Marial O, Stockler MR. Health status rated with the Medical Outcomes Study 36-Item Short-Form Health Survey after spinal cord injury. Arch Phys Med Rehabil. 2005;86(12):2290-2295.	10/10
179	Harper LA, Coleman JA, Perrin PB, et al. Comparison of mental health between individuals with spinal cord injury and able-bodied controls in Neiva, Colombia. Journal of rehabilitation research and development. 2014;51(1):127-136.	2/2
180	Harris EC, Barraclough BM, Grundy DJ, Bamford ES, Inskip HM. Attempted suicide and completed suicide in traumatic spinal cord injury. Case reports. Spinal Cord. 1996;34(12):752-753.	6/6
181	Harrison J, Glass CA, Owens RG, Soni BM. Factors associated with sexual functioning in women following spinal cord injury. Paraplegia. 1995;33(12):687-692.	10/10
82	Hart KA, Rintala DH, Fuhrer MJ. Educational interests of individuals with spinal cord injury living in the community: medical, sexuality, and wellness topics. Rehabil Nurs. 1996;21(2):82-90.	4/4
183	Hartkopp A, Bronnum-Hansen H, Seidenschnur AM, Biering-Sorensen F. Suicide in a spinal cord injured population: its relation to functional status. Arch Phys Med Rehabil. 1998;79(11):1356-1361.	10/10
184	Hartkopp PA, Bronnum-Hansen H, Seidenschnur AM, Biering-Sorensen F. [Survival and cause of death after traumatic spinal cord injury. A long-term epidemiological study]. Ugeskr Laeger. 1998;160(43):6207-6210.	8/8
85	Harvey JR, Bradley MB, Aronica MJ. Psychological effects of functional electrical stimulation: A review of the literature. Journal of Neurologic Rehabilitation. 1992;6(4):203-211.	11/11
86	Hassanijirdehi M, Khak M, Afshari-Mirak S, et al. Evaluation of pain and its effect on quality of life and functioning in men with spinal cord injury. Korean J Pain. 2015;28(2):129-136.	10/10
87	Hausmann LR, Myaskovsky L, Niyonkuru C, et al. Examining implicit bias of physicians who care for individuals with spinal cord injury: A pilot study and future directions. Journal of Spinal Cord Medicine. 2015;38(1):102-110.	4/4
188	Hayes RL, Potter CG, Hardin C. Counseling the client on wheels: A primer for mental health counselors new to spinal cord injury. Journal of Mental Health Counseling. 1995;17(1):18-30.	7/7
189	Heinemann A, Williams RT, Wilson CS, et al. Evaluating the psychometric properties of depression measures in persons with SCI and major depressive disorder. Archives of physical medicine and rehabilitation.94(10):e20-e21.	7/7
190	Heinemann AW, Doll MD, Armstrong KJ, Schnoll S, Yarkony GM. Substance use and receipt of treatment by persons with long-term spinal cord injuries. Arch Phys Med Rehabil. 1991;72(7):482-487.	10/10

Citation Number	Citation	Exclusion Code
191	Heinemann AW, Hawkins D. Substance abuse and medical complications following spinal cord injury. Rehabilitation Psychology. 1995;40(2):125-140.	10/10
192	Heinemann AW, Keen M, Donohue R, Schnoll S. Alcohol use by persons with recent spinal cord injury. Arch Phys Med Rehabil. 1988;69(8):619-624.	10/10
193	Heinemann AW, McGraw TE, Brandt MJ, Roth E, Dell'Oliver C. Prescription medication misuse among persons with spinal cord injuries. Int J Addict. 1992;27(3):301-316.	10/10
194	Heinemann AW, Schnoll S, Brandt M, Maltz R, Keen M. Toxicology screening in acute spinal cord injury. Alcohol Clin Exp Res. 1988;12(6):815-819.	4/4
195	Heinrich RK, Tate DG. Latent variable structure of the Brief Symptom Inventory in a sample of persons with spinal cord injuries. Rehabilitation Psychology. 1996;41(2):131-147.	2/2
196	Herrick SM, Elliott TR, Crow F. Social support and the prediction of health complications among persons with spinal cord injuries. Rehabilitation Psychology. 1994;39(4):231-250.	13/13
197	Heutink M, Post MW, Bongers-Janssen HM, et al. The CONECSI trial: results of a randomized controlled trial of a multidisciplinary cognitive behavioral program for coping with chronic neuropathic pain after spinal cord injury. Pain. 2012;153(1):120-128.	4/4
198	Heutink M, Post MW, Luthart P, Pfennings LE, Dijkstra CA, Lindeman E. A multidisciplinary cognitive behavioural programme for coping with chronic neuropathic pain following spinal cord injury: the protocol of the CONECSI trial. BMC neurology. 2010;10(96).	7/7
199	Heutink M, Post MWM, Bongers-Janssen HMH, et al. The conecsi trial: A randomized controlled trial of a multidisciplinary cognitive behavioral program for coping with chronic neuropathic spinal cord injury pain. Topics in spinal cord injury rehabilitation. 2011;16:29-30.	13/13
200	Hicken BL, Putzke JD, Richards JS. Bladder management and quality of life after spinal cord injury. Am J Phys Med Rehabil. 2001;80(12):916-922.	10/10
201	Hitzig SL, Tonack M, Campbell KA, et al. Secondary health complications in an aging Canadian spinal cord injury sample. Am J Phys Med Rehabil. 2008;87(7):545-555.	10/10
202	Hopkins MT. Patterns of self-destruction among the orthopedically disabled. Rehabilitation Research & Practice Review. 1971;3(1):5-16.	7/7
203	Horn SD, Smout RJ, DeJong G, et al. Association of various comorbidity measures with spinal cord injury rehabilitation outcomes. Arch Phys Med Rehabil. 2013;94(4 Suppl):S75-86.	4/4
204	Horner-Johnson W, Krahn GL, Suzuki R, et al. Differential performance of SF-36 items in healthy adults with and without functional limitations. Arch Phys Med Rehabil. 2010;91(4):570-575.	4/4
205	Horvat M, French R, Henschen K. A comparison of the psychological characteristics of male and female able-bodied and wheelchair athletes. Paraplegia. 1986;24(2):115-122.	4/4
206	Hossain MS, Rahman MA, Bowden JL, Quadir MM, Herbert RD, Harvey LA. Psychological and socioeconomic status, complications and quality of life in people with spinal cord injuries after discharge from hospital in Bangladesh: a cohort study. Spinal Cord. 2016;54(6):483-489.	2/2
207	Hough S, Stone MT, Buse DC. Dating and relationship psychoeducational group for veterans with spinal cord injury/dysfunction: A historical account of an initial clinical course. Sexuality and Disability. 2013;31(4):337-359.	4/4
208	Hough S. Depression after spinal cord injury and medication: The journey continues. Journal of Spinal Cord Medicine. 2014;37(3):246.	7/7
209	Houlihan BV, Jette A, Ni P, et al. Efficacy of "care call" telerehabilitation intervention for persons with spinal cord dysfunction: Randomized controlled trial. Archives of physical medicine and rehabilitation.92(10):1690.	7/7
210	Huang CY, Chen WK, Lu CY, et al. Mediating effects of social support and self- concept on depressive symptoms in adults with spinal cord injury. Spinal Cord. 2015;53(5):413-416.	4/4
211	Hui S-kA, Elliott TR, Shewchuk R, Rivera P. Communal behaviors and psychological adjustment of family caregivers and persons with spinal cord injury. Rehabilitation Psychology. 2007;52(1):113-119.	10/10
212	Hultling C, Giuliano F, Quirk F, Pena B, Mishra A, Smith MD. Quality of life in patients with spinal cord injury receiving Viagra (sildenafil citrate) for the treatment of erectile dysfunction. Spinal cord. 2000;38(6):363-370.	2/2
213	Hunter Revell SM. Symptom clusters in traumatic spinal cord injury: an exploratory literature review. J Neurosci Nurs. 2011;43(2):85-93.	4/4
214	Huston T, Gassaway J, Wilson C, Gordon S, Koval J, Schwebel A. The SCIRehab project: treatment time spent in SCI rehabilitation. Psychology treatment time during inpatient spinal cord injury rehabilitation. Journal of Spinal Cord Medicine. 2011;34(2):196-204.	4/4
215	Imai K, Kadowaki T, Aizawa Y. Standardized indices of mortality among persons with spinal cord injury: accelerated aging process. Ind Health. 2004;42(2):213-218.	4/4
216	Jacob KS, Zachariah K, Bhattacharji S. Depression in individuals with spinal cord injury: methodological issues. Paraplegia. 1995;33(7):377-380.	7/7
217	Jamous A, Kennedy P, Grey N. Psychological and emotional effects of the use of oral baclofen: A preliminary study. Paraplegia. 1994;32(5):349-353.	10/10

Continued

Citation Number	Citation	Exclusion Code
218	January AM, Zebracki K, Chlan KM, Vogel LC. Understanding post-traumatic growth following pediatric-onset spinal cord injury: the critical role of coping strategies for facilitating positive psychological outcomes. Dev Med Child Neurol. 2015;57(12):1143-1149.	13/13
219	Jensen MP, Hirsh AT, Molton IR, Bamer AM. Sleep problems in individuals with spinal cord injury: frequency and age effects. Rehabilitation Psychology. 2009;54(3):323-331.	4/4
220	Jensen MP, Hoffman AJ, Cardenas DD. Chronic pain in individuals with spinal cord injury: a survey and longitudinal study. Spinal Cord. 2005;43(12):704-712.	13/13
221	Jensen MP, Kuehn CM, Amtmann D, Cardenas DD. Symptom burden in persons with spinal cord injury. Arch Phys Med Rehabil. 2007;88(5):638-645.	13/13
222	Jensen MP, Moore MR, Bockow TB, Ehde DM, Engel JM. Psychosocial factors and adjustment to chronic pain in persons with physical disabilities: a systematic review. Arch Phys Med Rehabil. 2011;92(1):146-160.	11/11
223	Jiao J, Heyne MM, Lam CS. Acceptance of disability among Chinese individuals with spinal cord injuries: The effects of social support and depression. Psychology. 2012;3(9A):775-781.	13/13
224	Joerres SG, Bonifay RE, Hastings JE, Saltzstein RJ, Hayes TJ. Seasonal affective disorder in a spinal cord injury population. J Am Paraplegia Soc. 1992;15(2):66-70.	10/10
225	Johnson RL, Gerhart KA, McCray J, Menconi JC, Whiteneck GG. Secondary conditions following spinal cord injury in a population-based sample. Spinal Cord. 1998;36(1):45-50.	10/10
226	Johnstone B, Yoon DP. Relationships between the Brief Multidimensional Measure of Religiousness/ Spirituality and health outcomes for a heterogeneous rehabilitation population. Rehabilitation	10/10
227	Psychology. 2009;54(4):422-431. Judd FK, Brown DJ, Burrows GD. Depression, disease and disability: application to patients with traumatic spinal cord injury. Paraplegia. 1991;29(2):91-96.	11/11
228	Judd FK, Brown DJ. Psychiatric consultation in a spinal injuries unit. Australian and New Zealand Journal of Psychiatry. 1992;26(2):218-222.	10/10
229 230	Judd FK, Brown DJ. Psychiatry in the spinal injuries unit. Paraplegia. 1987;25(3):254-257. Judd FK, Brown DJ. Suicide following acute traumatic spinal cord injury. Paraplegia. 1992;30(3):173-177.	7/7 7/7
231	Judd FK, Burrows GD, Brown DJ. Depression following acute spinal cord injury. Paraplegia.	10/10
232	1986;24(6):358-363. Judd FK, Stone J, Webber JE, Brown DJ, Burrows G. Depression following spinal cord injury: A	10/10
233	prospective in-patient study. The British Journal of Psychiatry. 1989;154:668-671. Judd LL, Kunovac JL. Clinical characteristics and management of unipolar depressive disorders	6/6
234	comorbid with neurological disorders. Neuropsychobiology. 1998;37(2):84-87. Jurisic B, Marusic A. Suicidal ideation and behavior and some psychological correlates in physically	10/10
235	disabled motor-vehicle accident survivors. Crisis. 2009;30(1):34-38. Kalpakjian CZ, McCullumsmith CB, Fann JR, et al. Post-traumatic growth following spinal cord injury.	4/4
236	Journal of Spinal Cord Medicine. 2014;37(2):218-225. Kalpakjian CZ, Toussaint LL, Albright KJ, Bombardier CH, Krause JK, Tate DG. Patient health Questionnaire-9 in spinal cord injury: an examination of factor structure as related to gender. Journal of Spinal Cord Medicine. 2009;32(2):147-156.	4/4
237	Kannisto M, Sintonen H. Later health-related quality of life in adults who have sustained spinal cord injury in childhood. Spinal Cord. 1997;35(11):747-751.	10/10
238	Karimzadeh M, Salsabili N, Akbariasbagh F, Mehrsai A. The psychological impact of infertility in the male able bodied and spinal cord injured population. Sexuality and Disability. 2006;24(4):185-193.	10/10
239	Katz V, Gordon R, Iversen D, Myers SJ. Past history and degree of depression in paraplegic individuals. Paraplegia. 1978;16(1):8-14.	10/10
240	Kavussanu M, Ring C, Kavanagh J. Antisocial behavior, moral disengagement, empathy and negative emotion: A comparison between disabled and able-bodied athletes. Ethics & Behavior.	10/10
241	2015;25(4):297-306. Kawar JS, Sayegh HI, Shougah D, Tayseer F. Depression following spinal cord injury among	7/7
242	Jordanian patients. Arab Journal of Psychiatry. 1998;9(2):79-83. Kemp BJ, Kahan JS, Krause JS, Adkins RH, Nava G. Treatment of major depression in individuals	13/13
243	with spinal cord injury. Journal of Spinal Cord Medicine. 2004;27(1):22-28. Kemp BJ, Krause JS. Depression and life satisfaction among people ageing with post-polio and	4/4
244	spinal cord injury. Disabil Rehabil. 1999;21(5-6):241-249. Kemp BJ, Spungen AM, Adkins RH, Krause JS, Bauman WA. The relationships among serum lipid levels, adiposity, and depressive symptomatology in persons aging with spinal cord injury. Journal of Spinal Cord Madicine. 2000;00(4):243-269.	4/4
245	Spinal Cord Medicine. 2000;23(4):216-220. Kennedy P, Duff J, Evans M, Beedie A. Coping effectiveness training reduces depression and anxiety following traumatic spinal cord injuries. The British journal of clinical psychology / the British	13/13
246	Psychological Society. 2003;42(Pt 1):41-52. Kennedy P, Duff J. Post traumatic stress disorder and spinal cord injuries. Spinal Cord. 2001;39(1):1- 10.	11/11

Citation Number	Citation	Exclusion Code
247	Kennedy P, Evans M, Sandhu N. Psychological adjustment to spinal cord injury: the contribution of coping, hope and cognitive appraisals. Psychol Health Med. 2009;14(1):17-33.	13/13
248	Kennedy P, Evans MJ. Evaluation of post traumatic distress in the first 6 months following SCI. Spinal Cord. 2001;39(7):381-386.	2/2
249	Kennedy P, Kilvert A, Hasson L. A 21-year longitudinal analysis of impact, coping, and appraisals following spinal cord injury. Rehabilitation Psychology. 2016;61(1):92-101.	10/10
250	Kennedy P, Lowe R, Grey N, Short E. Traumatic spinal cord injury and psychological impact: A cross-sectional analysis of coping strategies. Br J Clin Psychol. 1995;34(4):627-639.	2/2
251	Kennedy P, Lude P, Elfstrom ML, Smithson E. Sense of coherence and psychological outcomes in people with spinal cord injury: appraisals and behavioural responses. Br J Health Psychol. 2010;15(Pt 3):611-621.	13/13
252	Kennedy P, Lude P, Elfstrom ML, Smithson EF. Psychological contributions to functional independence: a longitudinal investigation of spinal cord injury rehabilitation. Arch Phys Med Rehabil. 2011;92(4):597-602.	4/4
253	Kennedy P, Marsh N, Lowe R, Grey N, Short E, Rogers B. A longitudinal analysis of psychological impact and coping strategies following spinal cord injury. Br J Health Psychol. 2000;5(Part2):157-172.	13/13
254	Kennedy P, Rogers BA. Anxiety and depression after spinal cord injury: a longitudinal analysis. Arch Phys Med Rehabil. 2000;81(7):932-937.	13/13
255	Kennedy P, Taylor NM, Duff J. Characteristics Predicting Effective Outcomes After Coping Effectiveness Training for Patients With Spinal Cord Injuries. Journal of Clinical Psychology in Medical Settings. 2005;12(1):93-98.	10/10
256	Kennedy P. Coping effectively with spinal cord injuries: A group program: Therapist guide. Treatments that work. 2009.	2/2
257	Khazaeipour Z, Norouzi-Javidan A, Kaveh M, Khanzadeh Mehrabani F, Kazazi E, Emami-Razavi SH. Psychosocial outcomes following spinal cord injury in Iran. Journal of Spinal Cord Medicine. 2014;37(3):338-345.	2/2
258	Khazaeipour Z, Taheri-Otaghsara SM, Naghdi M. Depression Following Spinal Cord Injury: Its Relationship to Demographic and Socioeconomic Indicators. Topics in Spinal Cord Injury Rehabilitation. 2015;21(2):149-155.	2/2
259	Kilic SA, Dorstyn DS, Guiver NG. Examining factors that contribute to the process of resilience following spinal cord injury. Spinal Cord. 2013;51(7):553-557.	10/10
260	Kim SP, Davis SW, Sell GH. Amitriptyline in severely depressed spinal cord-injured patients: rapidity of response. Arch Phys Med Rehabil. 1977;58(4):157-161.	10/10
261	King C, Kennedy P. Coping effectiveness training for people with spinal cord injury: preliminary results of a controlled trial. The British journal of clinical psychology / the British Psychological Society. 1999;38(Pt 1):5-14.	10/10
262	Kirubakaran VR, Kumar VN, Powell BJ, Tyler AJ, Armatas PJ. Survey of alcohol and drug misuse in spinal cord injured veterans. Journal of Studies on Alcohol. 1986;47(3):223-227.	10/10
263	Kisala PA, Tulsky DS, Kalpakjian CZ, et al. Measuring anxiety after spinal cord injury: Development and psychometric characteristics of the SCI-QOL Anxiety item bank and linkage with GAD-7. Journal of Spinal Cord Medicine. 2015;38(3):315-325.	4/4
264	Kishi Y, Robinson RG, Forrester AW. Comparison between acute and delayed onset major depression after spinal cord injury. J Nerv Ment Dis. 1995;183(5):286-292.	10/10
265	Kishi Y, Robinson RG, Forrester AW. Prospective longitudinal study of depression following spinal cord injury. J Neuropsychiatry Clin Neurosci. 1994;6(3):237-244.	10/10
266	Kishi Y, Robinson RG, Kosier JT. Suicidal ideation among patients during the rehabilitation period after life-threatening physical illness. J Nerv Ment Dis. 2001;189(9):623-628.	5/5
267	Kishi Y, Robinson RG, Kosier JT. Suicidal ideation among patients with acute life- threatening physical illness: patients with stroke, traumatic brain injury, myocardial infarction, and spinal cord injury. Psychosomatics. 2001;42(5):382-390.	5/5
268	Kishi Y, Robinson RG. Suicidal plans following spinal cord injury: a six month study. J Neuropsychiatry Clin Neurosci. 1996;8(4):442-445.	13/13
269	Kivisild A, Sabre L, Tomberg T, et al. Health-related quality of life in patients with traumatic spinal cord injury in Estonia. Spinal Cord. 2014;52(7):570-575.	2/2
270	Kiwerski J, Weiss M, Chrostowska T. Analysis of mortality of patients after cervical spine trauma. Paraplegia. 1981;19(6):347-351.	4/4
271	Kiwerski JE, Krasuski M. Influence of alcohol intake on the course and consequences of spinal cord injury. International Journal of Rehabilitation Research. 1992;15(3):240-245.	4/4
272	Klyce DW, Bombardier CH, Davis TJ, et al. Distinguishing grief from depression during acute recovery from spinal cord injury. Arch Phys Med Rehabil. 2015;96(8):1419-1425.	4/4
273	Kohut RM, Seftel AD, Ducharme SH, Fogel BS, Bodner DR. Sexual and psychological aspects of rehabilitation after spinal cord injury. Psychiatric care of the medical patient. 2015:1504-1526.	6/6

Continued

NO. 1

Citation Number	Citation	Exclusion Code
274	Kolakowsky-Hayner SA, Gourley EV, 3rd, Kreutzer JS, Marwitz JH, Meade MA, Cifu DX. Post-injury substance abuse among persons with brain injury and persons with spinal cord injury. Brain Inj. 2002;16(7):583-592.	10/10
275	Kooijmans H, Post M, van der Woude L, de Groot S, Stam H, Bussmann J. Randomized controlled trial of a self-management intervention in persons with spinal cord injury: Design of the HABITS (Healthy Active Behavioural IntervenTion in SCI) study. Disability and Rehabilitation: An International, Multidisciplinary Journal. 2013;35(13):1111-1118.	7/7
276	Kortte KB, Gilbert M, Gorman P, Wegener ST. Positive psychological variables in the prediction of life satisfaction after spinal cord injury. Rehabilitation Psychology. 2010;55(1):40-47.	13/13
277	Kraft R, Dorstyn D. Psychosocial correlates of depression following spinal injury: A systematic review. Journal of Spinal Cord Medicine. 2015;38(5):571-583.	4/4
278	Kratz AL, Hirsh AT, Ehde DM, Jensen MP. Acceptance of pain in neurological disorders: associations with functioning and psychosocial well-being. Rehabilitation Psychology. 2013;58(1):1-9.	4/4
279	Krause J, Rohe DE. Personality and life adjustment after spinal cord injury: An exploratory study. Rehabilitation Psychology. 1998;43(2):118-130.	13/13
280	Krause JS, Bombardier C, Carter RE. Assessment of depressive symptoms during inpatient rehabilitation for spinal cord injury: Is there an underlying somatic factor when using the PHQ? Rehabilitation Psychology. 2008;53(4):513-520.	4/4
281	Krause JS, Broderick LE, Saladin LK, Broyles J. Racial disparities in health outcomes after spinal cord injury: mediating effects of education and income. Journal of Spinal Cord Medicine. 2006;29(1):17-25.	4/4
282	Krause JS, Brotherton SS, Morrisette DC, Newman SD, Karakostas TE. Does pain interference mediate the relationship of independence in ambulation with depressive symptoms after spinal cord injury? Rehabilitation Psychology. 2007;52(2):162-169.	4/4
283	Krause JS, Carter R, Zhai Y, Reed K. Psychologic factors and risk of mortality after spinal cord injury. Arch Phys Med Rehabil. 2009;90(4):628-633.	4/4
284	Krause JS, Coker J, Charlifue S, Whiteneck GG. Health behaviors among American Indians with spinal cord injury: comparison with data from the 1996 Behavioral Risk Factor Surveillance System. Arch Phys Med Rehabil. 1999;80(11):1435-1440.	10/10
285	Krause JS, Edles PA. Injury perceptions, hope for recovery, and psychological status after spinal cord injury. Rehabilitation Psychology. 2014;59(2):176-182.	2/2
286	Krause JS, Kjorsvig JM. Mortality after spinal cord injury: a four-year prospective study.[Erratum appears in Arch Phys Med Rehabil 1992 Aug;73(8):716]. Arch Phys Med Rehabil. 1992;73(6):558-563.	4/4
287	Krause JS, McArdle JJ, Pickelsimer E, Reed KS. A latent variable structural path model of health behaviors after spinal cord injury. Journal of Spinal Cord Medicine. 2009;32(2):162-174.	4/4
288	Krause JS, Reed KS, McArdle JJ. A structural analysis of health outcomes after spinal cord injury. Journal of Spinal Cord Medicine. 2010;33(1):22-32.	4/4
289	Krause JS, Reed KS, McArdle JJ. Factor structure and predictive validity of somatic and nonsomatic symptoms from the patient health questionnaire-9: a longitudinal study after spinal cord injury. Arch Phys Med Rehabil. 2010;91(8):1218-1224.	4/4
290	Krause JS, Reed KS, McArdle JJ. Prediction of somatic and non-somatic depressive symptoms between inpatient rehabilitation and follow-up. Spinal Cord. 2010;48(3):239-244.	10/10
291	Krause JS, Saunders LL, Bombardier C, Kalpakjian C. Confirmatory factor analysis of the Patient Health Questionnaire-9: a study of the participants from the spinal cord injury model systems. Pm R. 2011;3(6):533-540; quiz 540.	4/4
292	Krause JS, Vines CL, Farley TL, Sniezek J, Coker J. An exploratory study of pressure ulcers after spinal cord injury: relationship to protective behaviors and risk factors. Arch Phys Med Rehabil. 2001;82(1):107-113.	4/4
293	Krause JS, Zhai Y, Saunders LL, Carter RE. Risk of mortality after spinal cord injury: an 8-year prospective study. Arch Phys Med Rehabil. 2009;90(10):1708-1715.	13/13
294	Krause JS. Factors associated with risk for subsequent injuries after traumatic spinal cord injury. Arch Phys Med Rehabil. 2004;85(9):1503-1508.	4/4
295	Kreuter M, Sullivan M, Dahllof AG, Siosteen A. Partner relationships, functioning, mood and global quality of life in persons with spinal cord injury and traumatic brain injury. Spinal cord. 1998;36(4):252-261.	2/2
296	Kunz S, Joseph S, Geyh S, Peter C. Posttraumatic Growth and Adjustment to Spinal Cord Injury: Moderated by Posttraumatic Depreciation? Psychological Trauma: Theory, Research, Practice, and Policy. 2016:No Pagination Specified.	2/2
297	Kuptniratsaikul V, Chulakadabba S, Ratanavijitrasil S. An instrument for assessment of depression among spinal cord injury patients: comparison between the CES-D and TDI. J Med Assoc Thai. 2002;85(9):978-983.	2/2

Citation Number	Citation	Exclusion Code
298	Kuptniratsaikul V, Smerasuta O, Klomjaiyen P. The perceived handicap questionnaire: a self perceived handicap measurement in patients with spinal cord injury. J Med Assoc Thai. 2002;85(8):935-939.	10/10
299	Kuptniratsaikul V. Epidemiology of spinal cord injuries: a study in the Spinal Unit, Siriraj Hospital, Thailand, 1997-2000. J Med Assoc Thai. 2003;86(12):1116-1121.	10/10
300	Laatsch L, Shahani BT. The relationship between age, gender and psychological distress in rehabilitation inpatients. Disability and Rehabilitation: An International, Multidisciplinary Journal. 1996;18(12):604-608.	10/10
301	Lamid S, Chia JK, Kohli A, Cid E. Chronic pain in spinal cord injury: comparison between inpatients and outpatients. Arch Phys Med Rehabil. 1985;66(11):777-778.	4/4
302	Le J, Dorstyn D. Anxiety prevalence following spinal cord injury: a meta-analysis. Spinal Cord. 2016;54(8):626.	7/7
303	Levy LF, Makarawo S, Madzivire D, Bhebhe E, Verbeek N, Parry O. Problems, struggles and some success with spinal cord injury in Zimbabwe. Spinal Cord. 1998;36(3):213-218.	4/4
304	Lidal IB, Veenstra M, Hjeltnes N, Biering-Sorensen F. Health-related quality of life in persons with long-standing spinal cord injury.[Erratum appears in Spinal Cord. 2008 Nov;46(11):765]. Spinal Cord. 2008;46(11):710-715.	2/2
305	Liu S, Wang L, Cao C, Zhang J. The factor structure of posttraumatic stress disorder symptoms in patients with traumatic spinal cord injuries. Arch Psychiatr Nurs. 2014;28(5):301-304.	4/4
306	Livneh H, Antonak RF. Temporal structure of adaptation to disability. Rehabilitation Counseling Bulletin. 1991;34(4):298-319.	4/4
307	Livneh H, Martz E, Bodner T. Psychosocial Adaptation to Chronic Illness and Disability: A Preliminary Study of its Factorial Structure. Journal of Clinical Psychology in Medical Settings. 2006;13(3):251-261.	2/2
308	Livneh H, Martz E. Coping strategies and resources as predictors of psychosocial adaptation among people with spinal cord injury. Rehabilitation Psychology. 2014;59(3):329-339.	10/10
309	Livneh H, Martz E. Psychosocial adaptation to spinal cord injury as a function of time since injury. International Journal of Rehabilitation Research. 2003;26(3):191-200.	4/4
310	Livneh H, Martz E. The impact of perceptions of health control and coping modes on negative affect among individuals with spinal cord injuries. Journal of Clinical Psychology in Medical Settings. 2011;18(3):243-256.	10/10
311	Lohne V. The incomprehensible injuryinterpretations of patients' narratives concerning experiences with an acute and dramatic spinal cord injury. Scand J Caring Sci. 2009;23(1):67-75.	6/6
312	Lude P, Kennedy P, Evans M, Lude Y, Beedie A. Post traumatic distress symptoms following spinal cord injury: a comparative review of European samples. Spinal Cord. 2005;43(2):102-108.	10/10
313	Lundqvist C, Siosteen A, Blomstrand C, Lind B, Sullivan M. Spinal cord injuries. Clinical, functional, and emotional status. Spine. 1991;16(1):78-83.	4/4
314	Lynch KB, Bridle MJ. Construct validity of the Occupational Performance History Interview. Occupational Therapy Journal of Research. 1993;13(4):231-240.	2/2
315	Lyons JS, Larson DB, Burns BJ, Cope N, Wright S, Hammer JS. Psychiatric co-morbidities and patients with head and spinal cord trauma. Effects on acute hospital care. Gen Hosp Psychiatry. 1988;10(4):292-297.	4/4
316	MacDonald MR, Nielson WR, Cameron MG. Depression and activity patterns of spinal cord injured persons living in the community. Arch Phys Med Rehabil. 1987;68(6):339-343.	10/10
317 318	Mackenzie TB, Popkin MK. Suicide in the medical patient. Int J Psychiatry Med. 1987;17(1):3-22. MacKenzie TD, Kukolja T, House R, et al. A discharge panel at Denver Health, focused on complex	7/7 7/7
	patients, may have influenced decline in length-of-stay. Health Aff (Millwood). 2012;31(8):1786-1795.	·
319 320	Macleod AD. Self-neglect of spinal injured patients. Paraplegia. 1988;26(5):340-349. Macleod L, Macleod G. Control cognitions and psychological disturbance in people with contrasting	10/10 13/13
321	physically disabling conditions. Disabil Rehabil. 1998;20(12):448-456. Mann R, Schaefer C, Sadosky A, et al. Burden of spinal cord injury-related neuropathic pain in the United States: retrospective chart review and cross-sectional survey. Spinal Cord. 2013;51(7):564-570.	10/10
322	Martin Ginis KA, Jetha A, Mack DE, Hetz S. Physical activity and subjective well-being among people	11/11
323	with spinal cord injury: a meta-analysis. Spinal Cord. 2010;48(1):65-72. Martin Ginis KA, Jorgensen S, Stapleton J. Exercise and sport for persons with spinal cord injury. Pm R. 2012;4(11):894-900.	7/7
324	Martz E, Livneh H, Gontkovsky ST, Stokic DS. Psychosocial responses to spinal cord injury as predictors of pressure sores. International Journal of Clinical and Health Psychology. 2010;10(2):203-223.	4/4
325	Martz E, Livneh H, Priebe M, Wuermser LA, Ottomanelli L. Predictors of psychosocial adaptation	13/13
326	among people with spinal cord injury or disorder. Arch Phys Med Rehabil. 2005;86(6):1182-1192. Martz E, Livneh H. Death anxiety as a predictor of future time orientation among individuals with spinal cord injuries. Disabil Rehabil. 2003;25(18):1024-1032.	4/4

Citation Number	Citation	Exclusion Code		
327	Martz E. Death anxiety as a predictor of posttraumatic stress levels among individuals with spinal cord injuries. Death Stud. 2004;28(1):1-17.	2/2		
328	Martz E. Do post-traumatic stress symptoms predict reactions of adaptation to disability after a sudden-onset spinal cord injury? International Journal of Rehabilitation Research. 2004;27(3):185-194.	4/4		
329	Martz E. Do Reactions of Adaptation to Disability Influence the Fluctuation of Future Time Orientation Among Individuals with Spinal Cord Injuries? Rehabilitation Counseling Bulletin. 2004;47(2):86-95.	4/4		
330	Matsuda Y, Kubo T, Fujino Y, Matsuda S, Wada F, Sugita A. Relationship Between Depressive State and Treatment Characteristics of Acute Cervical Spinal Cord Injury in Japan. J Epidemiol. 2016;26(1):30-35.	2/2		
331	McAfee PC, Bohlman HH, Han JS, Salvagno RT. Comparison of nuclear magnetic resonance imaging and computed tomography in the diagnosis of upper cervical spinal cord compression. Spine. 1986;11(4):295-304.	6/6		
332	McColl MA, Rosenthal C. A model of resource needs of aging spinal cord injured men. Paraplegia. 1994;32(4):261-270.	13/13		
333	McColl MA, Skinner HA. Measuring psychological outcomes following rehabilitation. Can J Public Health. 1992;83 Suppl 2:S12-18.	4/4		
334	McColl MA, Skinner HA. Spinal cord injury and lifestyle health risks. Canadian Journal of Rehabilitation. 1995;9(2):69-82.	10/10		
335	McDermott S, Moran R, Platt T, Isaac T, Wood H, Dasari S. Risk for onset of health conditions among community-living adults with spinal cord and traumatic brain injuries. Primary Health Care Research and Development. 2007;8(1):36-43.	10/10		
336	McGowan MB, Roth S. Family functioning and functional independence in spinal cord injury adjustment. Paraplegia. 1987;25(4):357-365.			
337	McIntyre A, Cheung KY, Kwok C, Mehta S, Wolfe D, Teasell RW. Quality of life and bladder management post spinal cord injury: A systematic review. Applied Research in Quality of Life. 2014;9(4):1081-1096.	2/2		
338	McKinley WO, Kolakowsky SA, Kreutzer JS. Substance abuse, violence, and outcome after traumatic spinal cord injury. Am J Phys Med Rehabil. 1999;78(4):306-312.	4/4		
339	McKinley WO, Tewksbury MA, Godbout CJ. Comparison of medical complications following nontraumatic and traumatic spinal cord injury. Journal of Spinal Cord Medicine. 2002;25(2):88-93.			
340	Mehta S, McIntyre A, Dijkers M, Loh E, Teasell RW. Gabapentinoids are effective in decreasing neuropathic pain and other secondary outcomes after spinal cord injury: a meta-analysis. Arch Phys Med Rehabil. 2014;95(11):2180-2186.	11/11		
341	Mehta S, McIntyre A, Janzen S, Loh E, Teasell R, Spinal Cord Injury Rehabilitation Evidence T. Systematic Review of Pharmacologic Treatments of Pain After Spinal Cord Injury: An Update. Arch Phys Med Rehabil. 2016;97(8):1381-1391.e1381.	4/4		
342	Meyers AR, Bisbee A, Winter M. The "Boston model" of managed care and spinal cord injury: a cross-sectional study of the outcomes of risk-based, prepaid, managed care. Arch Phys Med Rehabil. 1999;80(11):1450-1456.	4/4		
343	Meyers AR, Feltin M, Master RJ, et al. Rehospitalization and spinal cord injury: cross-sectional survey of adults living independently. Arch Phys Med Rehabil. 1985;66(10):704-708.	4/4		
344	Migliorini C, Sinclair A, Brown D, Tonge B, New P. Prevalence of mood disturbance in Australian adults with chronic spinal cord injury. Intern Med J. 2015;45(10):1014-1019.	10/10		
345	Migliorini C, Tonge B. Reflecting on subjective well-being and spinal cord injury. Journal of Rehabilitation Medicine. 2009;41(6):445-450.	13/13		
346	Miller SM. The measurement of self-efficacy in persons with spinal cord injury: psychometric validation of the moorong self-efficacy scale. Disabil Rehabil. 2009;31(12):988-993.	4/4		
347	Miller WC, Anton HA, Townson AF. Measurement properties of the CESD scale among individuals with spinal cord injury. Spinal cord. 2008;46(4):287-292.	4/4		
348	Min JA, Lee CU, Hwang SI, et al. The moderation of resilience on the negative effect of pain on depression and post-traumatic growth in individuals with spinal cord injury. Disabil Rehabil. 2014;36(14):1196-1202.	10/10		
349	Molton IR, Stoelb BL, Jensen MP, Ehde DM, Raichle KA, Cardenas DD. Psychosocial factors and adjustment to chronic pain in spinal cord injury: replication and cross-validation. J Rehabil Res Dev. 2009;46(1):31-42.	4/4		
350	Moodley N, Pillay BJ. Post-traumatic stress disorder in patients with spinal-cord injuries. South African Journal of Psychology. 2013;43(2):182-197.	4/4		
351	Moore AD, Bombardier CH, Brown PB, Patterson DR. Coping and emotional attributions following spinal cord injury. International Journal of Rehabilitation Research. 1994;17(1):39-48.	13/13		
352	Mueller AD. Personality problems of the spinal cord injured. Journal of Consulting Psychology. 1950;14(3):189-192.	7/7		
353	Muller R, Cieza A, Geyh S. Rasch analysis of the Hospital Anxiety and Depression Scale in spinal cord injury. Rehabilitation Psychology. 2012;57(3):214-223.	4/4		

Citation Number	Citation	Exclusion Code
354	Muller R, Gertz KJ, Molton IR, et al. Effects of a Tailored Positive Psychology Intervention on Well-Being and Pain in Individuals with Chronic Pain and a Physical Disability. Clinical journal of pain. 2016;32(1):32-44.	4/4
355	Munce SE, Fehlings MG, Straus SE, et al. Views of people with traumatic spinal cord injury about the components of self-management programs and program delivery: a Canadian pilot study. BMC Neurology. 2014;14:209.	10/10
356	Muraki S, Tsunawake N, Hiramatsu S, Yamasaki M. The effect of frequency and mode of sports activity on the psychological status in tetraplegics and paraplegics. Spinal Cord. 2000;38(5):309-314.	4/4
357	Murray RF, Asghari A, Egorov DD, et al. Impact of spinal cord injury on self-perceived pre- and postmorbid cognitive, emotional and physical functioning. Spinal Cord. 2007;45(6):429-436.	10/10
358	Nagumo N. Depression occurring after delirium in traumatic spinal cord injury: A retrospective study. Seishin Igaku (Clinical Psychiatry). 1999;41(4):367-371.	8/8
359	Nestoros JN, Demers-Desrosiers LA, Dalicandro LA. Levels of anxiety and depression in spinal cordinjured patients. Psychosomatics. 1982;23(8):823-824, 827, 830.	10/10
360	Nicholls E, Lehan T, Plaza SL, et al. Factors influencing acceptance of disability in individuals with spinal cord injury in Neiva, Colombia, South America. Disabil Rehabil. 2012;34(13):1082-1088.	4/4
361	Nicholson Perry K, Nicholas MK, Middleton J, Siddall P. Psychological characteristics of people with spinal cord injury-related persisting pain referred to a tertiary pain management center. J Rehabil Res Dev. 2009;46(1):57-67.	10/10
362	Nicholson Perry K, Nicholas MK, Middleton J. Spinal cord injury-related pain in rehabilitation: a cross-sectional study of relationships with cognitions, mood and physical function. Eur J Pain. 2009;13(5):511-517.	13/13
363	Nicolais CJ, Perrin PB, Panyavin I, et al. Family dynamics and psychosocial functioning in children with SCI/D from Colombia, South America. Journal of Spinal Cord Medicine. 2016;39(1):58-66.	10/10
364	Nielsen MS. Post-traumatic stress disorder and emotional distress in persons with spinal cord lesion. Spinal Cord. 2003;41(5):296-302.	10/10
365	Nielson MS. Crisis support and coping as mediators of well-being in persons with spinal cord lesion. Journal of Clinical Psychology in Medical Settings. 2003;10(2):91-99.	10/10
366	Noonan VK, Kopec JA, Zhang H, Dvorak MF. Impact of associated conditions resulting from spinal cord injury on health status and quality of life in people with traumatic central cord syndrome. Arch Phys Med Rehabil. 2008;89(6):1074-1082.	10/10
367	Norrbrink Budh C, Hultling C, Lundeberg T. Quality of sleep in individuals with spinal cord injury: a comparison between patients with and without pain. Spinal Cord. 2005;43(2):85-95.	4/4
368	Norweg A, Jette AM, Houlihan B, Ni P, Boninger ML. Patterns, predictors, and associated benefits of driving a modified vehicle after spinal cord injury: findings from the National Spinal Cord Injury Model Systems. Arch Phys Med Rehabil. 2011;92(3):477-483.	3/3
369	Nott MT, Baguley IJ, Heriseanu R, et al. Effects of concomitant spinal cord injury and brain injury on medical and functional outcomes and community participation. Topics in Spinal Cord Injury Rehabilitation. 2014;20(3):225-235.	10/10
370	Osteraker AL, Levi R. Indicators of psychological distress in postacute spinal cord injured individuals. Spinal Cord. 2005;43(4):223-229.	10/10
371	Otis C, Marchand A, Courtois F. Peritraumatic dissociation as a mediator of peritraumatic distress and PTSD: a retrospective, cross-sectional study. J Trauma Dissociation. 2012;13(4):469-477.	10/10
372	Otis C, Marchand A, Courtois F. Risk factors for posttraumatic stress disorder in persons with spinal cord injury. Topics in Spinal Cord Injury Rehabilitation. 2012;18(3):253-263.	10/10
373	Overholser JC, Schubert DS, Foliart R, Frost F. Assessment of emotional distress following a spinal cord injury. Rehabilitation Psychology. 1993;38(3):187-198.	4/4
374	Paker N, Bugdayci D, Dere D, Altuncu Y. Comparison of the coping strategies, anxiety, and depression in a group of Turkish spinal cord injured patients and their family caregivers in a rehabilitation center. Eur J Phys Rehabil Med. 2011;47(4):595-600.	10/10
375	Paker N, Bugdayci D, Kesiktas N, Sahin M, Elfstrom ML. Reliability and validity of the Turkish version of spinal cord lesion-related coping strategies. Spinal Cord. 2014;52(5):383-387.	2/2
376	Paker N, Bugdayci D, Midik M, Celik B, Kesiktas N. Reliability of the Turkish version of the hospital anxiety and depression scale in the people with traumatic spinal cord injury. Neurorehabilitation. 2013;33(2):337-341.	2/2
377	Pang MY, Eng JJ, Lin KH, Tang PF, Hung C, Wang YH. Association of depression and pain interference with disease-management self-efficacy in community- dwelling individuals with spinal cord injury. Journal of Rehabilitation Medicine. 2009;41(13):1068-1073.	13/13
378	Paniak CE, Shore DL, Rourke BP, Finlayson MA, Moustacalis E. Long-term vocational functioning after severe closed head injury: a controlled study. Arch Clin Neuropsychol. 1992;7(6):529-540.	4/4
379	Paulsen P, French R, Sherrill C. Comparison of mood states of college able-bodied and wheelchair basketball players. Percept Mot Skills. 1991;73(2):396-398.	4/4
380	Pelletier PM, Alfano DP, Fink MP. Social support, locus of control and psychological health in family members following head or spinal cord injury. Appl Neuropsychol. 1994;1(1-2):38-44.	10/10

Citation Number	Citation	Exclusion Code
381	Perez M, Pilsecker C. Family therapy with spinal cord injured substance abusers. Soc Work Health Care. 1989;14(2):15-25.	7/7
382	Perez M, Pilsecker C. Group psychotherapy with spinal cord injured substance abusers. Paraplegia. 1994;32(3):188-192.	7/7
383	Perry KN, Nicholas MK, Middleton J. Spinal cord injury-related pain in rehabilitation: A cross-sectional study of relationships with cognitions, mood and physical function. Eur J Pain. 2009;13(5):511-517.	4/4
384	Perry KN, Nicholas MK, Middleton JW. Comparison of a pain management program with usual care in a pain management center for people with spinal cord injury-related chronic pain. Clinical Journal of Pain. 2010;26(3):206-216.	10/10
385	Peter C, Muller R, Cieza A, Geyh S. Psychological resources in spinal cord injury: a systematic literature review. Spinal Cord. 2012;50(3):188-201.	11/11
386	Pollard C, Kennedy P. A longitudinal analysis of emotional impact, coping strategies and post-traumatic psychological growth following spinal cord injury: a 10-year review. Br J Health Psychol. 2007;12(Pt 3):347-362.	13/13
387	Post MW, Adriaansen JJ, Charlifue S, Biering-Sorensen F, van Asbeck FW. Good validity of the international spinal cord injury quality of life basic data set. Spinal Cord. 2016;54(4):314-318.	6/6
388	Post MW, Gerritsen J, Diederiks JP, De Witte LP. Measuring health status of people who are wheelchair-dependent: Validity of the Sickness Impact Profile 68 and the Nottingham Health Profile. Disability and Rehabilitation: An International, Multidisciplinary Journal. 2001;23(6):245-253.	6/6
389	Post MW, van Leeuwen CM, van Koppenhagen CF, de Groot S. Validity of the Life Satisfaction questions, the Life Satisfaction Questionnaire, and the Satisfaction With Life Scale in persons with spinal cord injury. Arch Phys Med Rehabil. 2012;93(10):1832-1837.	6/6
390	Post MW, van Leeuwen CM. Psychosocial issues in spinal cord injury: a review. Spinal Cord. 2012;50(5):382-389.	11/11
391	Price GL, Kendall M, Amsters DI, Pershouse KJ. Perceived causes of change in function and quality of life for people with long duration spinal cord injury. Clinical Rehabilitation. 2004;18(2):164-171.	4/4
392	Price M. Causes of death in 11 of 227 patients with traumatic spinal cord injury over period of nine years. Paraplegia. 1973;11(3):217-220.	10/10
393	Putzke JD, Barrett JJ, Richards JS, DeVivo MJ. Age and spinal cord injury: an emphasis on outcomes among the elderly. Journal of Spinal Cord Medicine. 2003;26(1):37-44.	13/13
394	Putzke JD, Elliot TR, Richards J. Marital status and adjustment 1 year post-spinal-cord-injury. Journal of Clinical Psychology in Medical Settings. 2001;8(2):101-107.	4/4
395	Putzke JD, Hicken BL, Richards JS. Race: predictor versus proxy variable? Outcomes after spinal cord injury. Arch Phys Med Rehabil. 2002;83(11):1603-1611.	4/4
396	Putzke JD, Richards J, Dowler RN. The impact of pain in spinal cord injury: A case-control study. Rehabilitation Psychology. 2000;45(4):386-401.	10/10
397	Putzke JD, Richards JS, Hicken BL, DeVivo MJ. Interference due to pain following spinal cord injury: important predictors and impact on quality of life. Pain. 2002;100(3):231-242.	4/4
398	Quale AJ, Schanke AK, Froslie KF, Roise O. Severity of injury does not have any impact on posttraumatic stress symptoms in severely injured patients. Injury. 2009;40(5):498-505.	10/10
399	Quale AJ, Schanke AK. Resilience in the face of coping with a severe physical injury: a study of trajectories of adjustment in a rehabilitation setting. Rehabilitation Psychology. 2010;55(1):12-22.	10/10
400	Radnitz CL, Broderick CP, Perez-Strumolo L, et al. The prevalence of psychiatric disorders in veterans with spinal cord injury: a controlled comparison. J Nerv Ment Dis. 1996;184(7):431-433.	13/13
401	Radnitz CL, McGrath RE, Tirch DD, et al. Use of the Beck Depression Inventory in veterans with spinal cord injury. Rehabilitation Psychology. 1997;42(2):93-101.	13/13
402	Radnitz CL, Schlein IS, Hsu L. The effect of prior trauma exposure on the development of PTSD following spinal cord injury. J Anxiety Disord. 2000;14(3):313-324.	10/10
403	Rahnama P, Javidan AN, Saberi H, et al. Does religious coping and spirituality have a moderating role on depression and anxiety in patients with spinal cord injury? A study from Iran. Spinal Cord. 2015;53(12):870-874.	3/3
404	Raichle KA, Hanley M, Jensen MP, Cardenas DD. Cognitions, coping, and social environment predict adjustment to pain in spinal cord injury. Journal of Pain. 2007;8(9):718-729.	13/13
405	Reidy K, Caplan B. Causal factors in spinal cord injury: patients' evolving perceptions and association with depression. Arch Phys Med Rehabil. 1994;75(8):837-842.	10/10
406	Revell SM. Symptom clusters in traumatic spinal cord injury: An exploratory literature review. J Neurosci Nurs. 2011;43(2):85-93.	11/11
407	Richards JS, Bombardier CH, Wilson CS, et al. Efficacy of venlafaxine XR for the treatment of pain in patients with spinal cord injury and major depression: a randomized, controlled trial. Archives of physical medicine and rehabilitation. 2015;96(4):680-689.	4/4
408	Richards JS, Meredith RL, Nepomuceno C, Fine PR, Bennett G. Psycho-social aspects of chronic pain in spinal cord injury. Pain. 1980;8(3):355-366.	10/10

Continued

Citation Number	Citation	Exclusion Code
409	Richards JS, Osuna FJ, Jaworski TM, Novack TA, Leli DA, Boll TJ. The effectiveness of different methods of defining traumatic brain injury in predicting postdischarge adjustment in a spinal cord injury population. Arch Phys Med Rehabil. 1991;72(5):275-279.	4/4
410	Richards JS. Psychologic adjustment to spinal cord injury during first postdischarge year. Arch Phys Med Rehabil. 1986;67(6):362-365.	10/10
411	Richardson EJ, Richards J. Factor structure of the PHQ-9 screen for depression across time since injury among persons with spinal cord injury. Rehabilitation Psychology. 2008;53(2):243-249.	4/4
412	Rintala DH, Young ME, Hart KA, Clearman RR, Fuhrer MJ. Social support and the well-being of persons with spinal cord injury living in the community. Rehabilitation Psychology. 1992;37(3):155-163.	13/13
413	Rintala DH, Young ME, Hart KA, Fuhrer MJ. The relationship between the extent of reciprocity with social supporters and measures of depressive symptomatology, impairment, disability, and handicap in persons with spinal cord injury. Rehabilitation Psychology. 1994;39(1):15-27.	13/13
414	Rintala DH. Predictive validity of social support relative to psychological well-being in men with spinal cord injury. Rehabilitation Psychology. 2013;58(4):422-428.	4/4
415	Riordan A, Kelly EH, Klaas SJ, Vogel LC. Psychosocial outcomes among youth with spinal cord injury by neurological impairment. Journal of Spinal Cord Medicine. 2015;38(1):76-83.	4/4
416	Rish BL, Dilustro JF, Salazar AM, Schwab KA, Brown HR. Spinal cord injury: a 25-year morbidity and mortality study. Mil Med. 1997;162(2):141-148.	10/10
417	Robinson-Whelen S, Taylor HB, Hughes RB, Wenzel L, Nosek MA. Depression and depression treatment in women with spinal cord injury. Topics in Spinal Cord Injury Rehabilitation. 2014;20(1):23-31.	10/10
418	Rodrigues D, Tran Y, Wijesuriya N, Guest R, Middleton J, Craig A. Pain intensity and its association with negative mood States in patients with spinal cord injury. Pain Ther. 2013;2(2):113-119.	4/4
419	Rosenberg DE, Bombardier CH, Artherholt S, Jensen MP, Motl RW. Self-reported depression and physical activity in adults with mobility impairments. Arch Phys Med Rehabil. 2013;94(4):731-736.	11/11
420	Roth E, Davidoff G, Thomas P, et al. A controlled study of neuropsychological deficits in acute spinal cord injury patients. Paraplegia. 1989;27(6):480-489.	2/2
421	Rothstein JL, Levy E, Fecher R, Gordon SK, Bauman WA. Drug use and abuse in an urban veteran spinal cord injured population. J Am Paraplegia Soc. 1992;15(4):217-220.	10/10
422	Rowland JL, White GW, Wyatt DA. Analysis of an intervention to reduce or prevent secondary conditions for people with spinal cord injuries. Journal of clinical psychology in medical settings. 2006;13(3):263-271.	6/6
423	Russell HF, January AM, Kelly EH, Mulcahey MJ, Betz RR, Vogel LC. Patterns of coping strategy use and relationships with psychosocial health in adolescents with spinal cord injury. J Pediatr Psychol. 2015;40(5):535-543.	2/2
424	Saadat S, Javadi M, Divshali BS, et al. Health-related quality of life among individuals with long- standing spinal cord injury: a comparative study of veterans and non-veterans. BMC Public Health. 2010;10:6.	4/4
425	Sabre L, Rekand T, Asser T, Korv J. Mortality and causes of death after traumatic spinal cord injury in Estonia. Journal of Spinal Cord Medicine. 2013;36(6):687-694.	2/2
426	Sadosky A, Parsons B, Emir B, Nieshoff EC. Pain relief and functional improvement in patients with neuropathic pain associated with spinal cord injury: an exploratory analysis of pregabalin clinical trials. J Pain Res. 2016;9:405-416.	4/4
427	Saikkonen J, Karppi P, Huusko TM, Dahlberg A, Makinen J, Uutela T. Life situation of spinal cord- injured persons in Central Finland. Spinal Cord. 2004;42(8):459-465.	13/13
428	Salinas FA, Lugo LH, Garcia HI. Efficacy of early treatment with carbamazepine in prevention of neuropathic pain in patients with spinal cord injury. American journal of physical medicine &	11/11
429	rehabilitation / Association of Academic Physiatrists. 2012;91(12):1020-1027. Salonia A, Munarriz RM, Naspro R, et al. Women's sexual dysfunction: a pathophysiological review.	7/7
430	BJU Int. 2004;93(8):1156-1164. Salter JE, Smith SD, Ethans KD. Positive and negative affect in individuals with spinal cord injuries. Spinal Cord. 2013;51(3):252-256.	10/10
431	Sammallahti P, Kannisto M, Aalberg V. Psychological defenses and psychiatric symp- toms in adults with pediatric spinal cord injuries. Spinal Cord. 1996;34(11):669-672.	4/4
432	Saulino M, Averna JF. Evaluation and Management of SCI-Associated Pain. Curr Pain Headache Rep. 2016;20(9):53.	6/6
433	Saunders LL, Krause JS, Carpenter MJ, Saladin M. Risk behaviors related to cigarette smoking among persons with spinal cord injury. Nicotine Tob Res. 2014;16(2):224-230.	2/2
434	Schandler SL, Cohen MJ, Vulpe M. Problem solving and coping strategies in persons with spinal cord injury who have and do not have a family history of alcoholism. Journal of Spinal Cord Medicine. 1996;19(2):78-86.	10/10
435	Schnoll SH, Heinemann AW, Doll M, Armstrong KJ. Substance use and receipt of treatment in persons with recent spinal cord injuries. NIDA Res Monogr. 1989;95:426-427.	6/6

Citation Number	Citation	Exclusion Code
436	Schulz R, Decker S. Long-term adjustment to physical disability: the role of social support, perceived control, and self-blame. J Pers Soc Psychol. 1985;48(5):1162-1172.	13/13
437	Scivoletto G, Petrelli A, Di Lucente L, Castellano V. Psychological investigation of spinal cord injury patients. Spinal Cord. 1997;35(8):516-520.	2/2
438	Shadish WR, Hickman D, Arrick M. Psychological problems of spinal cord injury patients: Emotional distress as a function of time and locus of control. Journal of Consulting and Clinical Psychology. 1981;49(2):297.	7/7
439	Shin JC, Goo HR, Yu SJ, Kim DH, Yoon SY. Depression and Quality of Life in Patients within the First 6 Months after the Spinal Cord Injury. Ann. 2012;36(1):119-125.	10/10
440	Shin JI, Chae JH, Min JA, et al. Resilience as a possible predictor for psychological distress in chronic spinal cord injured patients living in the community. Ann. 2012;36(6):815-820.	10/10
441	Shnek ZM, Foley FW, LaRocca NG, et al. Helplessness, self-efficacy, cognitive distortions, and depression in multiple sclerosis and spinal cord injury. Ann Behav Med. 1997;19(3):287-294.	13/13
442	Skinner AL, Armstrong KJ, Rich J. Depression and spinal cord injury: A review of diagnostic methods for depression, 1985 to 2000. Rehabilitation Counseling Bulletin. 2003;46(3):174-175.	7/7
443	Smedema SM, Ebener D. Substance abuse and psychosocial adaptation to physical disability: analysis of the literature and future directions. Disabil Rehabil. 2010;32(16):1311-1319.	11/11
444	Smith BM, Weaver FM, Ullrich PM. Prevalence of depression diagnoses and use of antidepressant medications by veterans with spinal cord injury. Am J Phys Med Rehabil. 2007;86(8):662-671.	13/13
445	Smith TF, Russell HF, Kelly EH, Mulcahey MJ, Betz RR, Vogel LC. Examination and measurement of coping among adolescents with spinal cord injury. Spinal Cord. 2013;51(9):710-714.	2/2
446	Soden RJ, Walsh J, Middleton JW, Craven ML, Rutkowski SB, Yeo JD. Causes of death after spinal cord injury. Spinal Cord. 2000;38(10):604-610.	10/10
447	Somasundaram O, Balakrishnan S, Ravindran OS, Shanmugasundaram TK. A psychological study of spinal cord injured patients involved in the Madras Paraplegia Project. Paraplegia. 1992;30(11):799-802.	10/10
448	Stroud MW, Bombardier CH, Dyer JR, Rimmele CT, Esselman PC. Preinjury alcohol and drug use among persons with spinal cord injury: implications for rehabilitation. Journal of Spinal Cord Medicine. 2011;34(5):461-472.	13/13
449	Stroud MW, Turner JA, Jensen MP, Cardenas DD. Partner responses to pain behaviors are associated with depression and activity interference among persons with chronic pain and spinal cord injury. Journal of Pain. 2006;7(2):91-99.	13/13
450	Summers JD, Rapoff MA, Varghese G, Porter K, Palmer RE. Psychosocial factors in chronic spinal cord injury pain. Pain. 1991;47(2):183-189.	13/13
451	Tamplin J, Baker FA, Grocke D, Berlowitz DJ. Thematic analysis of the experience of group music therapy for people with chronic quadriplegia. Topics in Spinal Cord Injury Rehabilitation. 2014;20(3):236-247.	10/10
452	Tasiemski T, Kennedy P, Gardner BP, Blaikley RA. Athletic Identity and Sports Participation in People With Spinal Cord Injury. Adapt Phys Act Q. 2004;21(4):364-378.	13/13
453	Tasiemski T, Kennedy P, Gardner BP, Taylor N. The association of sports and physical recreation with life satisfaction in a community sample of people with spinal cord injuries. Neurorehabilitation. 2005;20(4):253-265.	4/4
454	Tate DG, Forchheimer M, Bombardier CH, Heinemann AW, Neumann HD, Fann JR. Differences in quality of life outcomes among depressed spinal cord injury trial participants. Archives of physical medicine and rehabilitation. 2015;96(2):340-348.	4/4
455	Tate DG, Forchheimer M, Maynard F, Davidoff G, Dijkers M. Comparing two measures of depression in spinal cord injury. Rehabilitation Psychology. 1993;38(1):53-61.	13/13
156	Tate DG, Forchheimer MB, Karana-Zebari D, Chiodo AE, Kendall Thomas JY. Depression and pain among inpatients with spinal cord injury and spinal cord disease: differences in symptoms and neurological function. Disabil Rehabil. 2013;35(14):1204-1212.	4/4
457	Tate DG, Heinrich RK, Maynard F, Buckelew SP. Moderator-variable effect on the Brief Symptom Inventory test-item endorsements of spinal cord injury patients. Paraplegia. 1994;32(7):473-479.	4/4
458	Tate DG, Kalpakjian C, Chiodo A, et al. Treatment of mild to moderate depression in persons with SCI: A randomized clinical trial. Archives of physical medicine and rehabilitation.93(10):E9.	7/7
159	Tate DG, Kewman DG, Maynard F. The Brief Symptom Inventory: Measuring psychological distress in spinal cord injury. Rehabilitation Psychology. 1990;35(4):211-216.	4/4
460	Tate DG. Alcohol use among spinal cord-injured patients. Am J Phys Med Rehabil. 1993;72(4):192-195.	13/13
461	Tavakoli SA, Kavian M, Bakhsh SC, et al. Is Level of Injury a Determinant of Quality of Life Among Individuals with Spinal Cord Injury? A Tertiary Rehabilitation Center Report. Oman med. 2016;31(2):112-116.	2/2
462	Tawashy AE, Eng JJ, Lin KH, Tang PF, Hung C. Physical activity is related to lower levels of pain, fatigue and depression in individuals with spinal-cord injury: a correlational study. Spinal Cord. 2009;47(4):301-306.	10/10

Continued

Citation Number	Citation	Exclusion Code	
463	Tee JW, Chan CH, Gruen RL, et al. Early predictors of health-related quality of life outcomes in polytrauma patients with spine injuries: a level 1 trauma center study. Global spine j. 2014;4(1):21-32.	4/4	
464	Temple RO, Elliott TR. Personality disorder characteristics and adjustment following spinal cord injury. Topics in Spinal Cord Injury Rehabilitation. 2000;6(1):54-65.	10/10	
465	Tetrault M, Courtois F. Use of psychoactive substances in persons with spinal cord injury: a literature review. Ann Phys Rehabil Med. 2014;57(9-10):684-695.	11/11	
466	Tirch D, Radnitz CL, Bauman WA. Depression and spinal cord injury: a monozygotic twin study. Journal of Spinal Cord Medicine. 1999;22(4):284-286.	11/11	
467	Tomey KM, Chen DM, Wang X, Braunschweig CL. Dietary intake and nutritional status of urban community-dwelling men with paraplegia. Arch Phys Med Rehabil. 2005;86(4):664-671.	4/4	
468	Tramonti F, Gerini A, Stampacchia G. Relationship quality and perceived social support in persons with spinal cord injury. Spinal Cord. 2015;53(2):120-124.	10/10	
469	Tulsky DS, Kisala PA, Kalpakjian CZ, et al. Measuring depression after spinal cord injury: Development and psychometric characteristics of the SCI-QOL Depression item bank and linkage with PHQ-9. Journal of Spinal Cord Medicine. 2015;38(3):335-346.	6/6	
470	Turner AP, Bombardier CH, Rimmele CT. A typology of alcohol use patterns among persons with recent traumatic brain injury or spinal cord injury: implications for treatment matching. Arch Phys Med Rehabil. 2003;84(3):358-364.	4/4	
471	Turner JA, Jensen MP, Warms CA, Cardenas DD. Catastrophizing is associated with pain intensity, psychological distress, and pain-related disability among individuals with chronic pain after spinal cord injury. Pain. 2002;98(1-2):127-134.	4/4	
472	Ullrich P, Jensen M, Loeser J, Cardenas D. Catastrophizing mediates associations between pain severity, psychological distress, and functional disability among persons with spinal cord injury. Rehabilitation Psychology. 2007;52(4):390-398.	2/2	
473	Ullrich PM, Smith BM, Poggensee L, et al. Pain and post-traumatic stress disorder symptoms during inpatient rehabilitation among operation enduring freedom/ operation iraqi freedom veterans with spinal cord injury. Arch Phys Med Rehabil. 2013;94(1):80-85.	13/13	
474	Valtonen K, Karlsson AK, Alaranta H, Viikari-Juntura E. Work participation among persons with traumatic spinal cord injury and meningomyelocele1. Journal of Rehabilitation Medicine. 2006;38(3):192-200.	4/4	
475	van Leeuwen CM, Edelaar-Peeters Y, Peter C, Stiggelbout AM, Post MW. Psychological factors and mental health in persons with spinal cord injury: An exploration of change or stability. Journal of Rehabilitation Medicine. 2015;47(6):531-537.	10/10	
476	van Leeuwen CM, Hoekstra T, van Koppenhagen CF, de Groot S, Post MW. Trajectories and predictors of the course of mental health after spinal cord injury. Arch Phys Med Rehabil. 2012;93(12):2170-2176.	2/2	
477	van Leeuwen CM, Post MW, Westers P, et al. Relationships between activities, participation, personal factors, mental health, and life satisfaction in persons with spinal cord injury. Arch Phys Med Rehabil. 2012;93(1):82-89.	2/2	
478	van Leeuwen CM, van der Woude LH, Post MW. Validity of the mental health subscale of the SF-36 in persons with spinal cord injury. Spinal Cord. 2012;50(9):707-710.	6/6	
479	Virseda-Chamorro M, Salinas-Casado J, Lopez-Garcia-Moreno AM, Cobo-Cuenca AI, Esteban-Fuertes M. Sexual dysfunction in men with spinal cord injury: a case-control study. Int J Impot Res. 2013;25(4):133-137.	10/10	
480	Vissers M, van den Berg-Emons R, Sluis T, Bergen M, Stam H, Bussmann H. Barriers to and facilitators of everyday physical activity in persons with a spinal cord injury after discharge from the rehabilitation centre. Journal of Rehabilitation Medicine. 2008;40(6):461-467.	4/4	
481	Weaver FM, LaVela SL, Miskevics S, Clemmons N, Janke EA, Spring B. Smoking behavior and readiness to change in male veterans with spinal cord injuries. Rehabilitation Psychology. 2007;52(3):304-310.	4/4	
482	Weaver FM, Smith B, LaVela SL, et al. Smoking behavior and delivery of evidence-based care for veterans with spinal cord injuries and disorders. Journal of Spinal Cord Medicine. 2011;34(1):35-45.	13/13	
483	Weeks DL, Greer CL, Bray BS, Schwartz CR, White JR, Jr. Association of antidepressant medication therapy with inpatient rehabilitation outcomes for stroke, traumatic brain injury, or traumatic spinal cord injury. Arch Phys Med Rehabil. 2011;92(5):683-695.	4/4	
484	Weingardt KR, Hsu J, Dunn ME. Brief screening for psychological and substance abuse disorders in veterans with long-term spinal cord injury. Rehabilitation Psychology. 2001;46(3):271-278.	10/10	
485	Westgren N, Levi R. Quality of life and traumatic spinal cord injury. Arch Phys Med Rehabil. 1998;79(11):1433-1439.	4/4	
486	White B, Driver S, Warren AM. Resilience and indicators of adjustment during rehabilitation from a spinal cord injury. Rehabilitation Psychology. 2010;55(1):23-32.	10/10	
487	Widerstrom-Noga EG, Felipe-Cuervo E, Yezierski RP. Chronic pain after spinal injury: interference with sleep and daily activities. Arch Phys Med Rehabil. 2001;82(11):1571-1577.	4/4	

Citation Number	Citation	Exclusion Code
488	Wilcox NE, Stauffer ES. Follow-up of 423 consecutive patients admitted to the Spinal Cord Centre, Rancho Los Amigos Hospital, 1 January to 31 December 1967. Paraplegia. 1972;10(2):115-122.	10/10
489	Williams RT, Heinemann AW, Bode RK, Wilson CS, Fann JR, Tate DG. Improving measurement properties of the Patient Health Questionnaire-9 with rating scale analysis. Rehabilitation Psychology. 2009;54(2):198-203.	4/4
490	Williamson ML, Elliott TR. Major depressive disorder and factorial dimensions among individuals with recent-onset spinal cord injury. Rehabilitation Psychology. 2013;58(1):10-17.	4/4
491	Wollaars MM, Post MW, van Asbeck FW, Brand N. Spinal cord injury pain: the influence of psychologic factors and impact on quality of life. Clinical Journal of Pain. 2007;23(5):383-391.	13/13
492	Wong AW, Heinemann AW, Wilson CS, et al. Predictors of participation enfranchisement after spinal cord injury: the mediating role of depression and moderating role of demographic and injury characteristics. Arch Phys Med Rehabil. 2014;95(6):1106-1113.	4/4
493	Wood-Dauphinee S, Exner G, Bostanci B, et al. Quality of life in patients with spinal cord injurybasic issues, assessment, and recommendations. Restor Neurol Neurosci. 2002;20(3-4):135-149.	7/7
494	Wool RN, Siegel D, Fine PR. Task performance in spinal cord injury: effect of helplessness training. Arch Phys Med Rehabil. 1980;61(7):321-325.	10/10
495	Woolrich RA, Kennedy P, Tasiemski T. A preliminary psychometric evaluation of the Hospital Anxiety and Depression Scale (HADS) in 963 people living with a spinal cord injury. Psychol Health Med. 2006;11(1):80-90.	13/13
496	Young ME, Rintala DH, Rossi CD, Hart KA, Fuhrer MJ. Alcohol and marijuana use in a community-based sample of persons with spinal cord injury. Arch Phys Med Rehabil. 1995;76(6):525-532.	10/10
497	Znoj HJ, Lude P. Regulation of emotion and psychological symptoms in people with spinal cord injury. Swiss Journal of Psychology / Schweizerische Zeitschrift fur Psychologie / Revue Suisse de Psychologie. 2002;61(4):203-210.	4/4

Appendix D. Evidence Tables

Table D-1. Risk of bias of systematic reviews

Author, Year	Was an "a priori" design provided?	Was there duplicate study selection?	Was there duplicate data extraction?	Was a comprehensive literature search performed?	Was the status of publication (i.e., grey literature) used as an inclusion criterion?	Was inclusion/ exclusion criteria determined a priori?
Boldt, 2014	Yes	Yes	Yes	Yes	No	Yes
Craig, 2009	Yes	Unclear	Unclear	Yes	Unclear	Yes
Dorstyn, 2011 (CBT)	Yes	Unclear	Unclear	Yes	Unclear	Yes
Dorstyn, 2013 (telehealth)	Yes	Unclear	Unclear	Yes	Unclear	Yes
Jones, 2015	Yes	Unclear	No	Yes	No	Unclear
Kalpakjian, 2009	Yes	Yes	Partial	Yes	No	Partial
Le, 2016	Yes	Yes	Yes	Yes	Yes	Yes
Lusilla- Palacios, 2015	Yes	Unclear	Yes	Yes	No	Yes
Mehta, 2014	Yes	Yes	Yes	Yes	Yes	Yes
Muller, 2012	Yes	Yes	Yes	Yes	Unclear	Yes
Perkes, 2014	Yes	Yes	Yes	Yes	Unclear	Yes
Sakakibara, 2009	Yes	Yes	Yes	Yes	No	Yes
van Gorp, 2014	Yes	Unclear	Unclear	Yes	Unclear	Yes
van Leeuwen, 2012	Yes	Yes	Unclear	Yes	Unclear	Yes
Williams, 2015	Yes	Yes	No	Yes	Unclear	Yes

Author, Year	Was a list of included studies provided?	Was a list of excluded studies provided?	Were the characteristics of the included studies provided?	Was the scientific quality of the included studies assessed and documented?	Was the scientific quality of the included studies used appropriately in formulating conclusions?
Boldt, 2014	Yes	Yes	Yes	Yes	Yes
Craig, 2009	Yes	No	Yes	No	No
Dorstyn, 2011 (CBT)	Yes	No	Yes	Yes	Yes
Dorstyn, 2013 (telehealth)	Yes	No	Yes	Yes	Yes
Jones, 2015	Yes	No	Yes	No	No
Kalpakjian, 2009	Yes	No	Yes	No	No
Le, 2016	Yes	No	Yes	Unclear	Unclear
Lusilla- Palacios, 2015	Yes	No	Yes	No	Unclear
Mehta, 2014	Yes	No	Yes	Yes	Yes
Muller, 2012	No	No	Yes	Yes	Yes
Perkes, 2014	Yes	No	Yes	Yes	Yes
Sakakibara, 2009	Yes	No	Yes	Yes	Yes
van Gorp, 2014	Yes	No	Yes	Yes	Yes
van Leeuwen, 2012	Yes	No	Yes	Yes	Yes
Williams, 2015	Yes	No	Yes	Unclear	Unclear

Author, Year	Were the methods used to combine the findings of studies appropriate? Do the qualitative syntheses and Strength of Evidence ratings appropriately consider factors such as precision, consistency, and methodological limitations?	Was the likelihood of publication bias assessed?	Were conflicts of interest described for the systematic review authors?	Were conflicts of interest described for authors of the primary studies?	Risk of Bias
Boldt, 2014	Yes	No	Yes	No	Low
Craig, 2009	Unclear	No	No	No	High
Dorstyn, 2011 (CBT)	Yes	Unclear	No	No	Medium
Dorstyn, 2013 (telehealth)	Yes	Unclear	Yes	No	Medium
Jones, 2015	Unclear	No	Yes	No	High
Kalpakjian, 2009	Yes	No	No	No	Medium
Le, 2016	Yes	Yes	Yes	No	Low
Lusilla- Palacios, 2015	No	No	Yes	No	Medium

Author, Year	the D	re the methods used to combine findings of studies appropriate? of the qualitative syntheses and Strength of Evidence ratings opriately consider factors such as precision, consistency, and methodological limitations?	Was the likelihood of publication bias assessed?	Were conflicts of interest described for the systematic review authors?	Were conflicts of interest described for authors of the primary studies?	Risk of Bias
Mehta, 2014	Yes		No	Yes	No	Medium
Muller, 2012	Yes		No	Yes	No	Medium
Perkes, 2014	Yes		Yes	Yes	No	Low
Sakakibara, 2009	Yes		No	No	No	Medium
van Gorp, 2014	Yes		No	Yes	No	Medium
van Leeuwen, 2012	Yes		No	Yes	No	Medium
Williams, 2015	Yes		Yes	No	No	Medium

Table D-2. Risk of bias of randomized controlled trials

Author, Year	Random- ization adequate?	Allocation concealment adequate?	Groups similar at baseline?	Outcome assessors masked?	Care provider masked?	Patient masked?	Intention- to-treat analysis	Acceptable level of overall attrition (<20%)	Acceptable level of differential attrition (<10%)	Risk of Bias
Cardenas, 2002	Yes	Unclear	Yes	Yes	Yes	Yes	Yes	No	Yes	Medium
Cardenas, 2013	Yes	Yes	Yes	Unclear	Yes	Yes	Yesv	Yes	Yes	Low
Diego, 2002 Fann, 2015 Bombardier, 2014 Ruhe, 2005 Williams, 2016	No Yes	Unclear Yes	Unclear Yes	Yes Yes	No Yes	No Yes	Unclear Yes	Unclear Yes	Unclear Yes	High Low
Hicks, 2003 Martin- Ginnis, 2003	No	Unclear	Yes	Unclear	No	No	Yes	Yes	Yes	High
Houlihan, 2011 Houlihan, 2013 Mercier, 2015	Unclear	Unclear	No	Unclear	No	No	Yes	Yes	Yes	Medium
Nooijen, 2016	Yes	Unclear	Yes	Yes	No	No	Yes	No	Yes	Medium
Norrbrink, 2009	Unclear	Unclear	No	Unclear	Yes	Yes	Yes	Yes	Yes	High
Rintala, 2007	Yes	Unclear	Yes	Unclear	Unclear, double- blind	Unclear, double- blind	Yes	No	Yes	Medium
Siddall, 2006	Yes	Yes	Yes	Unclear	Unclear, double- blind	Unclear, double- blind	Yes	No	Yes	Medium
Yilmaz, 2014	Unclear	Unclear	Unclear	Unclear	Yes	Unclear, double- blind	Unclear	No	Yes	High

Table D-3. Risk of bias of observational studies

Author, Year	Non- biased selection?	Acceptable overall loss to follow-up or differential loss to follow-up?	Outcomes pre- specified and defined?	Ascertainment techniques ade- quately described?	Non-biased and adequate ascertainment methods?	Statistical analysis of potential confounders?	Adequate duration of follow-up?	Overall risk of bias rating (High, Medium, Low)
Ahoniemi, 2011	Yes	Unclear	Yes	Yes	Yes	Unclear	Yes	Medium
Bradley, 1994	No	No; Yes	Yes	Yes	Yes	No	Yes	High
Hartkopp, 1997	Yes	Unclear	Unclear	Yes	Yes	Unclear	Yes	Medium
Kalpakjian, 2006	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Medium
Kemp, 1999 "depression among african american	Unclear	Yes	Yes	Yes	Yes	Unclear	Yes	High
Kennedy, 2006 (psychosocial activity) Krause, 1999	Unclear	Yes	Yes	Yes	Yes	No	Yes	High
Krause, 2000 "health outcomes"	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Medium
Krause, 2009	Yes	Yes	Yes	Yes	Yes	Unclear	Yes	Medium
Sweeney, 1982	No	Yes	Unclear	Yes	Unclear	No	Yes	High
Ullrich, 2014	Yes	Yes	Yes	Yes	Yes	Unclear	Yes	Medium

Table D-4. Risk of bias of observational studies of risk factors

Author, Year	The study sample adequately represents the population of interest?	The study data available (i.e., participants not lost to followup) adequately represent the study sample?	The prognostic factor is measured in a similar way for all participants?	The outcome of interest is measured in a similar way for all participants?	Important potential confounding factors are appropriately accounted for?	The statistical analysis is appropriate, and all primary outcome are reported?	Risk of Bias
Abdul-Sattar, 2014	Yes	Yes	Yes	Yes	Yes	Yes	Medium
Alschuler, 2013	Yes	Yes	Yes	Yes	Yes	Yes	Medium
Arango-Lasprilla, 2011	Yes	Yes	Yes	Yes	Yes	Yes	Medium
Bailey, 2012	Yes	Yes	Yes	Yes	Yes	Yes	Medium
Bannerjea, 2009	Yes	Yes	Yes	Yes	Yes	Yes	Medium
Binks, 1997	Unclear	Unclear	Unclear	Unclear	Unclear	Unclear	High
Bombardier, 2016	Yes	Yes	Yes	Yes	Yes	Yes	Medium
Burns, 2009 Burns, 2010	Yes	Unclear	Yes	Yes	Yes	Yes	Medium
Cao, 2014 (Rehab Psychology)	Unclear	Yes	Yes	Yes	Yes	Yes	Medium
Cao 2014b (Archives of)	Yes	Yes	Yes	Yes	Yes	Yes	Medium
Catalano, 2011	Unclear	Unclear	Yes	Yes	Yes	Unclear	Medium
Charlifue, 1991	Yes	Yes	Yes	Yes	Yes	Yes	Medium
Cuff, 2014	Yes	Yes	Yes	Yes	Yes	Yes	Medium
Cragg, 2015	Unclear	Unclear	Yes	Yes	Yes	Yes	High
Craig, 2015	Unclear	Unclear	Yes	Yes	Yes	Yes	High
Danner, 2000	Yes (VA)	Yes	Yes	Yes	Yes	Yes	Medium
Daverat, 1995	Yes	Yes	Yes	Yes	Yes	Yes	Medium
Dodd, 2015	Yes	Yes	Yes	Yes	Yes	Yes	Medium
Dryden, 2005	Yes	Yes	Yes	Yes	Yes	Yes	Medium
Fann, 2013	Yes	Yes	Yes	Yes	Yes	Yes	Medium
Geyh, 2013	Yes	Yes	Yes	Yes	Yes	Yes	Medium
Gioia, 2006	Yes	Yes	Yes	Yes	Yes	Yes	Medium
Hartoonian, 2014	Yes	Yes	Yes	Yes	Yes	Yes	Medium

Table D-4. Continued

Author, Year	The study sample adequately represents the population of interest?	The study data available (i.e., participants not lost to followup) adequately represent the study sample?	The prognostic factor is measured in a similar way for all participants?	The outcome of interest is measured in a similar way for all participants?	Important potential confounding factors are appropriately accounted for?	The statistical analysis is appropriate, and all primary outcome are reported?	Risk of Bias
Hassanpour,	Yes	Yes	Yes	Yes	Yes	Yes	Medium
2012 Hatcher, 2009	Unclear	Yes	Yes	Yes	Yes	Yes	High
Hoffman, 2011	Yes	Yes	Yes	Yes	Yes	Yes	Medium
Holicky, 1999	Yes	Yes	Yes	Yes	Yes	Yes	Medium
Huang, 2014	Yes	Yes	Yes	Yes	Yes	Yes	Medium
Hwang, 2014	Yes	Yes	Yes	Yes	Yes	Yes	Medium
January, 2014	Yes	Yes	Yes	Yes	Yes	Yes	Medium
(Archives of Phys	162	165	162	165	165	162	Medium
Med and Rehab)	Yes	Yes	Yes	Yes	Yes	Yes	Medium
January, 2015 (Rehabilitation Psych)	res	res	res	res	res	res	Medium
Jensen, 2014	Yes	Yes	Yes	Yes	Yes	Yes	Medium
Kalpakjian, 2011	Yes	Yes	Yes	Yes	Yes	Yes	Medium
Kemp, 2014	Yes	Yes	Yes	Yes	Yes	Yes	Medium
Krause, 2010	Yes	Yes	Yes	Yes	Yes	Yes	Medium
(arch phys med rehabil)							
LaVela, 2012	Unclear	Unclear	Yes	Yes	Yes	Yes	High
Lidal, 2013	Unclear	Unclear	Yes	Yes	Yes	Yes	High
Lin, 2009	Unclear	Unclear	Yes	Yes	Yes	Yes	High
Liu, 2010	Unclear	Yes	Yes	Yes	Yes	Yes	Medium
Lundqvist, 1997	Unclear	Yes	Yes	Yes	Yes	Yes	Medium
Martz, 2005 (rehab psych)	Unclear	Yes	Yes	Yes	Yes	Yes	Medium
McCullumsmith, 2015	Yes	Yes	Yes	Yes	Yes	Yes	Medium
Migliorini, 2008	Yes	Yes	Yes	Yes	Yes	Yes	Medium
Migliorini, 2009	Yes	Yes	Yes	Yes	Yes	Yes	Medium
Monin, 2014	Yes	Yes	Yes	Yes	Yes	Yes	Medium
Muller, 2015	Yes	Yes	Yes	Yes	Yes	Yes	Medium
Nam, 2013	Yes	Yes	Yes	Yes	Yes	Yes	Medium
Nelson, 2010	Yes	Yes	Yes	Yes	Yes	Yes	Medium
Nielsen, 2003	Yes	Yes	Yes	Yes	Yes	Yes	High
O'Shea, 2014	Yes	Yes	Yes	Yes	Yes	Yes	Medium
Oh, 2006	Yes	Yes	Yes	Yes	Yes	Yes	Medium
Pakpour, 2016	Yes	Yes	Yes	Yes	Yes	Yes	Medium
Peter, 2015	Yes	Yes	Yes	Yes	Yes	Yes	Medium
Riggins, 2011	Yes	Yes	Yes	Yes	Yes	Yes	Medium
Rintala, 2005	Yes	Yes	Yes	Yes	Yes	Yes	Medium
Saunders, 2011	Yes	Unclear	Yes	Yes	Yes	Yes	Medium
Saunders, 2013							
Saunders, 2012	Yes	Yes	Yes	Yes	Yes	Yes	Medium
Schandler, 1995	Yes	Yes	Yes	Yes	Yes	Yes	Medium
Schonenberg, 2014	Yes	Yes	Yes	Yes	Yes	Yes	Medium
Tasiemski, 2011	Yes	Yes	Yes	Yes	Yes	Yes	Medium
Tate, 1994	Yes	Yes	Yes	Yes	Yes	Yes	Medium
Tate, 2004	Yes	Yes	Yes	Yes	Yes	Yes	Medium
Tsai, 2014	Yes	Yes	Yes	Yes	Yes	Yes	Medium
Ullrich, 2013	Yes	Yes	Yes	Yes	Yes	Yes	Medium
(rehabilitation psychology)							
Ullrich, 2014	Yes	Yes	Yes	Yes	Yes	Yes	High
Williams, 2014	Yes	Yes	Yes	Yes	Yes	Yes	Medium