

Supplemental data 4. K-seq protocol

1- First synthesis

1 μ l of each forward short-primers (5 μ M each)
2 μ l dNTPs (25mM)
4,5 μ l klenow buffer 10X
200-600 ng genomic DNA
H₂O until 45 μ l final volume

Reaction:

4 min 95°C
5 min 37°C—add 1 μ l Klenow (5U/ μ l Klenow Fragment (3'→5' exo))
20 min 75°C

2- First Exonuclease digestion

+ 2 μ l Exonuclease I (E. coli) 20U/ μ l

Reaction:

60 min 37°C
20 min 80°C

3- Second synthesis

+1 μ l each reverse short-primers (5 μ M each)
+0.5 buffer Klenow 10X
+ H₂O to 4 μ l final volume

Reaction:

4 min 95°C
5 min 37°C. In this step add 1 μ l Klenow (5U/ μ l)
20 min 75°C

4- Second Exonuclease digestion

+ 1 μ l Exonuclease I (E. coli) 20U/ μ l

Reaction:

60 min 37°C
20 min 80°C

5- PCR

15 μ l reaction 4
25 μ l Taq NEB 2X (M0270L Taq 2X Master MixNEB)
1 μ l IDT-NXT adapter i7 (2.5 μ M)
1 μ l IDT-NXT adapter i5 (2.5 μ M)
8 μ l H₂O

Reaction:

1 min 95°C
X cycles (usually between 10 and 15 cycles. Adjust cycle number to avoid high molecular weight band smear in agarose gel electrophoresis)
30s 95°C
20s 62°C
30s 68°C
Last extension step. 5 min 68°C