

Note to readers with disabilities: *EHP* strives to ensure that all journal content is accessible to all readers. However, some figures and Supplemental Material published in *EHP* articles may not conform to [508 standards](#) due to the complexity of the information being presented. If you need assistance accessing journal content, please contact ehp508@niehs.nih.gov. Our staff will work with you to assess and meet your accessibility needs within 3 working days.

Supplemental Material

Effect of Epidermal Growth Factor Treatment and Polychlorinated Biphenyl Exposure in a Dietary-Exposure Mouse Model of Steatohepatitis

Josiah E. Hardesty, Banrida Wahlang, Russell A. Prough, Kim Z. Head, Daniel Wilkey, Michael Merchant, Hongxue Shi, Jian Jin, and Matthew C. Cave

Table of Contents

Table S1. PCB Congener Composition of Aroclor1260 (Adapted from ATSDR CDC).

Table S2. Taqman Probes.

Table S3. Antibodies.

Table S4. Outliers for each measure and figure.

Table S5. p-values.

Table S6. Numerical Data.

Figure S1. Effects of acute EGF and chronic EGF and Aroclor1260 exposure on hepatic EGFR signaling.

Figure S2. Effects of chronic EGF and Aroclor 1260 exposure on fasted glucose levels at 6 and 12 hours.

Figure S3. Effects of chronic EGF and Aroclor 1260 exposure on hepatic FXR and AhR expression and target genes.

Figure S4. Effect of chronic EGF and A1260 exposure on hepatic phosphoproteomic targets.

Additional File- Excel Document