Supplemental table 1: General characteristics of clinical characteristics of excluded versus included data

••	Included Data	Excluded Data	••
•••	n (%)	n (%)	p-value ^A
Outcome Total	215937 (%)	9514 (%)	
Age Median (IQR)	23.0 (19.0, 28.0)	23.0 (20.0, 29.0)	< 0.001
Under 20	58759 (27.2%)	2323 (24.4%)	
20-29	112721 (52·2%)	5013 (52.7%)	
30 or Older	44452 (20.6%)	2178 (22.9%)	••
Unknown	5 (<1%)	0 (0.0%)	
Self-Identified Race	••		< 0.001
White	67076 (31·1%)	2027 (21.3%)	••
Black	25731 (11.9%)	873 (9.2%)	
Mixed or Indigenous	106860 (49.5%)	4051 (42.6%)	••
Unknown	16270 (7.5%)	2563 (26.9%)	••
Education			< 0.001
Some Primary			
School	66053 (30.6%)	2004 (21·1%)	
Some Secondary			
School	50862 (23.6%)	1294 (13.6%)	
HS Graduate or			
Higher	40134 (18.6%)	1047 (11.0%)	
Unknown	58888 (27.3%)	5169 (54·3%)	
Population Density	••		< 0.001
Urban	67076 (31·1%)	2027 (21.3%)	
Rural or Periurban	25731 (11.9%)	873 (9.2%)	
Unknown	106860 (49.5%)	4051 (42.6%)	••

A. p- value obtained from Kruskal-Wallis tests for continuous variables, and χ^2 tests for categorical variables

Supplemental table 2: Clinical characteristics of excluded versus included data

	Included Data	Excluded Data	
	n (%)	n (%)	p-value ^A
Outcome Total	215937	9514	••
Prenatal Care	••	••	< 0.001
No Prenatal Care	16899 (7.8%)	2760 (29.0%)	••
Received Prenatal Care	199038 (92.2%)	6754 (71.0%)	
Syphilis Titer	••	••	< 0.001
<16	116782 (54·1%)	5989 (62.9%)	
≥16	99155 (45.9%)	3525 (37·1%)	
Term of Diagnosis	••	••	< 0.001
First Term	68836 (31.9%)	1703 (17.9%)	
Second Term	63079 (29·2%)	1646 (17.3%)	
Third Term or Later	83808 (38.8%)	6154 (64.7%)	••
Unknown	214 (0.1%)	11 (0.1%)	••

A. p-value obtained from Kruskal-Wallis tests for continuous variables, and χ^2 tests for categorical variables