




## **Supporting Information**

### **Supplementary methods and results**

**This appendix was part of the submitted manuscript and has been peer reviewed.  
It is posted as supplied by the authors.**

Appendix to: Darley DR, Dore GJ, Cysique L, et al. Persistent symptoms up to four months after community and hospital-managed SARS-CoV-2 infection. *Med J Aust* 2021; doi: 10.5694/mja2.50963.

**Figure 1. Participant selection and assessment**



ARDS = acute respiratory distress syndrome; COVID-19 = coronavirus disease 2019; ICU = intensive care unit; PCR = polymerase chain reaction; SARS-CoV-2 = severe acute respiratory syndrome coronavirus 2.

**Table 1. Definitions of ADAPT cohort sub-populations**

Sub-population	
Mild	Managed in the community; minor, largely upper respiratory tract viral symptoms, including sore throat, rhinorrhoea, headache, and anosmia/ageusia
Moderate	Managed in the community with fever/chills <b>AND</b> one of the following organ-localising symptoms, <b>OR</b> at least two of the following organ-localising symptoms: cough, haemoptysis, shortness of breath, chest pain, nausea/vomiting, diarrhea, altered consciousness/confusion
Severe	Required inpatient care (wards or intensive care unit)

**Table 2. Patient characteristics at the time of COVID-19 presentation**

Patient characteristic	All patients	Community	Hospitalised
Number of patients	78	69	9
Age at diagnosis (years), mean (SD)	47 (16)	45 (15)	62 (8.6)
Sex (women)	27	27	0
Ethnic background (European)	65	57	8
Epidemiology			
- Overseas travel	39	35	4
- Known close contact	38	32	6
- Presence in a high-risk setting	15	12	3
Comorbid conditions			
- None	37	36	1
- Hypertension	14	12	2
- Diabetes mellitus	7	3	4
- Chronic cardiac disease	8	5	3
- Chronic lung disease	1	1	0
- Asthma	9	8	1
- Chronic kidney disease	3	2	1
- Cancer	7	6	1
- Immune disorders	3	3	0
- Obesity	3	2	1
- Obstructive sleep apnoea	1	1	0
- Psychiatric diagnosis	6	6	0
Smoking (ever)	29	21	8
Highest education level			
- University	61	54	7
- High school	14	13	1
ICU admission for ARDS	2	—	2

ARDS = acute respiratory distress syndrome; ICU = intensive care unit.