

Supporting Information

**Production, characterization, and antioxidant activities of
exopolysaccharide extracted from spent media wastewater after
Leuconostoc mensenteroides WiKim32 fermentation**

In Seong Choi^{a,b}, Seung Hee Ko^{a,b}, Mo Eun Lee^a, Ho Myeong Kim^a, Jung Eun Yang^a, Seul-Gi Jeong^a, Kwang Ho Lee^c, Ji Yoon Chang^a, Jin-Cheol Kim^d and Hae Woong Park^{a,*}


^aR&D Division, World Institute of Kimchi, Gwangju, 61755, Republic of Korea

^bPublic CMO for Microbial – based Vaccine, Hwasun-gun, Jeollanam-do, 58141, Republic of Korea


^cCenter for Research Facilities, Chonnam National University, Gwangju, 61186, Republic of Korea

^dDepartment of Agricultural Chemistry, Institute of Environmentally Friendly Agriculture, College of Agriculture and Life Science, Chonnam National University, Gwangju 61186, Republic of Korea

* Corresponding author at: Advanced Process Technology Fermentation Research Group, R&D Division, World Institute of Kimchi, Gwangju 61755, Republic of Korea



Supplementary Figure S1. The UV spectra of EPS in the range 200-900 nm.



Supplementary Figure S2. FTIR spectrum of the commercial dextran.