

Supplemental Table 1. Signs and symptoms present at time of transplantation

Pt Number	Onset (mo)	Age of Transplant (mo)	RIC	Global Delay	Abnormal gait	Abnormal muscle tone	Reflux	Constipation	Feeding Difficulty	Tracking Difficulty	Abnormal Pupillary Response	Staring Spells	Clonus	Abnormal DTR	Seizure
1	N/A	13	0	0	0	0	0	0	0	0	0	0	0	0	0
2	N/A	40	0	0	0	0	0	0	0	0	0	0	0	0	0
3	N/A	1	0	0	0	0	0	0	0	0	0	0	0	0	0
4	N/A	3	0	0	0	0	0	0	0	0	0	0	0	0	0
5	N/A	4	0	0	0	0	0	0	0	0	0	0	0	0	0
6	10	14	0	1	0	1	0	0	0	0	0	0	0	1	0
7	15	34	0	0	1	1	0	0	0	0	1	0	0	0	0
8	10.5	15	0	1	0	1	0	0	1	0	1	0	0	1	0
9	12	23	0	1	0	1	0	0	0	0	0	0	0	0	1
10	8	11	0	1	0	1	0	0	1	0	0	0	1	1	0
11	12	21	0	0	0	1	1	0	1	0	0	0	0	0	1
12	10	18	0	1	0	1	0	1	0	0	0	0	0	1	0
13	31	38	0	0	1	1	0	0	0	0	0	1	1	1	0
14	28	32	1	0	1	1	0	1	0	0	0	0	1	1	0
15	17	19	1	1	0	1	1	1	0	0	0	0	0	0	0
16	7.5	12	0	0	0	1	1	1	1	0	0	1	0	1	0
17	24	28	1	0	1	0	0	1	0	0	0	1	0	1	0
18	6	8	1	0	0	1	1	1	1	1	0	0	1	1	0
19	35	44	1	1	1	1	0	1	0	0	0	1	0	0	0

Supplemental Table 1 displays selected signs and symptoms that were present in patients at the time of transplantation. While data exists on a more diverse and inclusive set of signs and symptoms, the ones selected here represent the most frequent and pertinent found in this cohort of patients. A “1” indicates that the sign/symptom was present; a “0” indicates that the sign/symptom was not present. Patients with N/A were asymptomatic at the time of transplantation.