Dimeric mimetic of BDNF loop 4 promotes survival of serum-deprived cell through TrkB-dependent apoptosis suppression

L.F. Zainullina §, Yu.V. Vakhitova §*, A.Yu. Lusta, T.A. Gudasheva, S.B. Seredenin

Federal State Budgetary Institution «Research Zakusov Institute of Pharmacology»,

Moscow, Russia

§authors contributed equally to this work.

*Correspondence: Vakhitova Yulia, vakhitova@academpharm.ru; Department of Pharmacogenetics, «Research Zakusov Institute of Pharmacology»,

Baltiyskaya str., 8, Moscow, 125315, Russia

Supplementary Info File

Data availability statement: All the original data acquisition of western blots is included in Supplementary Info file.


Keywords: BDNF, low weight mimetics, TrkB signaling, neuroprotection, BAD, GSK-3β

Supplementary Figure Legend


Supplementary Figure 1. (**A**) MEK1 inhibitor (PD98059) had no effect on Akt pathway. SH-SY5Y cells in serum-free culture medium («0% FBS») with or without BDNF (~ 1 nmol) or GSB-106 (100 nmol) were also pre incubated with or without PD98059 (50 μmol, 60 min). Blots were first probed with anti-phosphorylated pan-Akt (pAkt1/2/3) antibody and then reprobed with anti-pan-Akt antibody, anti-a-tubulin antibody. The figure shows data from one independent experiment (n=3; the original blots are shown in the supplementary information file).

(**B**) PI3K inhibitor (LY294002) had no effect on MAPK pathway. SH-SY5Y cells in serum-free culture medium («0% FBS») with or without BDNF (~ 1 nmol) or GSB-106 (100 nmol) were also pre incubated with or without LY290042 (50 μmol, 60 min). Blots were first probed with anti-phosphorylated Erk (pErk) antibody and then reprobed with anti-Erk antibody, anti-a-tubulin antibody. The figure shows data from one independent experiment (n=3; the original blots are shown in the supplementary information file).


Zainullina et al., Supplementary Figure 1









Original scans of blots, corresponding to figure 1B







Original scans of blots, corresponding to figure 3A




Original scans of blots, corresponding to figure 3C




Original scans of blots, corresponding to figure 3E







Original scans of blots, corresponding to figure 4A




Original scans of blots, corresponding to figure 4D




Original scans of blots, corresponding to figure 4E




Original scans of blots, corresponding to figure 4F




Zainullina et al., Figure 5 as submitted




Original scans of blots, corresponding to figure 5A




Original scans of blots, corresponding to figure 5C




Original scans of blots, corresponding to figure 5C (continued)




Original scans of blots, corresponding to figure 5D



Zainullina et al., Figure 6 as submitted




Original scans of blots, corresponding to figure 6A and 6E



anti-α-Tubulin (1:1000, #2125, Cell Signaling Technology)

anti-α-Tubulin (1:1000, #2125, Cell Signaling Technology)

Zainullina et al., Supplementary Figure 1 as submitted



Zainullina et al., Supplementary Figure 1

Original scans of blots, corresponding to Supplementary Figure 1A and 1B

