

Note to readers with disabilities: *EHP* strives to ensure that all journal content is accessible to all readers. However, some figures and Supplemental Material published in *EHP* articles may not conform to [508 standards](#) due to the complexity of the information being presented. If you need assistance accessing journal content, please contact ehp508@niehs.nih.gov. Our staff will work with you to assess and meet your accessibility needs within 3 working days.

Supplemental Material

Fluoride in Drinking Water, Diet, and Urine in Relation to Bone Mineral Density and Fracture Incidence in Postmenopausal Women

Emilie Helte, Carolina Donat Vargas, Maria Kippler, Alicja Wolk, Karl Michaëlsson, and Agneta Åkesson

Table of Contents

Table S1. Cross-sectional mean differences in BMD (β coefficients (95% CI), g/cm²) across tertiles of urinary fluoride (mg/g creatinine) and dietary fluoride (mg/d) with BMD at the lumbar spine and femoral neck among women with approximately constant drinking water fluoride concentrations from 1982 to baseline.

Table S2. Hazard ratios of osteoporotic fractures, excluding vertebral fractures, and corresponding 95% confidence intervals by tertiles of urinary fluoride (mg/g creatinine) and dietary fluoride (mg/d).

Table S3. Multivariable adjusted hazard ratios of total, osteoporotic and hip fractures and corresponding 95% confidence intervals by tertiles of urinary fluoride (mg/g creatinine) and dietary fluoride (mg/d), respectively. Multivariable models are further adjusted for baseline bone mineral density (BMD).

Table S4. Cross-sectional mean differences in BMD (β coefficients (95% CI), g/cm²) across tertiles of urinary fluoride (mg/g creatinine) and dietary fluoride (mg/d) with BMD at the lumbar spine and femoral neck, among women of the Swedish Mammography Cohort – Clinical, including individuals excluded from main analysis because of missing bone mineral density data on either hip or spine.

Table S5. Hazard ratios of total, osteoporotic and hip fractures and corresponding 95% confidence intervals by tertiles of urinary fluoride (mg/g creatinine) and dietary fluoride (mg/d), respectively, among women of the Swedish Mammography Cohort – Clinical, including individuals excluded from main analysis because of missing bone mineral density data on either hip or spine.

Table S6. Cross-sectional mean differences in Bone mineral density (BMD)(β coefficients (95% CI), g/cm^2) across tertiles of urinary fluoride (mg/g creatinine) with BMD at the lumbar spine and femoral neck among women of the Swedish Mammography Cohort – Clinical with complete exposure data.

Table S7. Hazard ratios of total, osteoporotic and hip fractures and corresponding 95% confidence intervals by tertiles of urinary fluoride (mg/g creatinine) among women of the Swedish Mammography Cohort – Clinical with complete exposure data.

Table S8. Cross-sectional mean differences in Bone mineral density (BMD)(β coefficients (95% CI), g/cm^2) across tertiles of urinary fluoride (mg/g creatinine) with BMD at the lumbar spine and femoral neck among women of the Swedish Mammography Cohort – Clinical with urinary creatinine concentrations ranging between 0.3 g/L and 3.0 g/L .

Table S9. Hazard ratios of total, osteoporotic and hip fractures and corresponding 95% confidence intervals by tertiles of urinary fluoride (mg/g creatinine) among women of the Swedish Mammography Cohort – Clinical with urinary creatinine concentrations ranging between 0.3 g/L and 3.0 g/L .