

Supplemental Table 1 Primer sequences used for qRT-PCR.

Gene	Forward primer (5' to 3')	Reverse primer (5' to 3')
<i>Col1α1</i>	GTGCTCCTGGTATTGCTGGT	GGCTCCTCGTTTTTCCTTCTT
<i>Col3α1</i>	GGGTTTCCCTGGTCCTAAAG	CCTGGTTTCCCATTTTCTCC
<i>Col6α1</i>	GATGAGGGTGAAGTGGGAGA	CAGCACGAAGAGGATGTCAA
<i>Fibronectin</i>	ACACGGTTTCCCATTACGCCAT	AATGACCACTGCCAAAGCCCAA
<i>LOX</i>	AGCTTGCTTTGTGGCCTTCA	CCACAGCATGGACGAATTCA
<i>F4/80</i>	CTTTGGCTATGGGCTTCCAGTC	GCAAGGAGGACAGAGTTTATCGTG
<i>TNFα</i>	CAACCTCCTCTCTGCCGTCAA	TGACTCCAAAGTAGACCTGCCC
<i>IL-1β</i>	GCTACCTGTGTCTTTCCCGT	CGTCACACACCAGCAGGTTA
<i>IL-6</i>	AGTTGCCTTCTTGGGACTGA	ACAGGTCTGTTGGGAGTGGT
<i>MCP-1</i>	CCCAATGAGTAGGCTGGAGA	TCTGGACCCATTCCTTCTTG
<i>HIF1α</i>	CAAGATCTCGGCGAAGCAA	GGTGAGCCTCATAACAGAAGCTTT
<i>PPAR γ</i>	ATCTTAACTGCCGGATCCAC	CAAACCTGATGGCATTGTGAG
<i>β-Actin</i>	CGTGAAAAGATGACCCAGATCA	CAGCCTGGATGGCTACGTACA