

S1 Table. Epidemiologic data of selected non-malignant primary brain and other CNS tumors

	Vestibular schwannoma	Pituitary adenoma	WHO Gr I Meningioma	WHO Gr II Meningioma
10 Year total	12,491	21,472	39,363	1,591
Median age	54.0	49.0	61.0	58.0
Male sex	5,603	8,634	8,350	616
Female sex	6,888	12,838	31,013	975
All age (95% CI) ^{a)}	2.31 (2.26-2.35)	4.00 (3.94-4.05)	7.52 (7.45-7.60)	0.30 (0.29-0.32)
0-14 yr (95% CI) ^{b)}	0.10 (0.08-0.13)	0.30 (0.26-0.34)	0.07 (0.05-0.09)	0.02 (0.01-0.03)
15-39 yr (95% CI) ^{b)}	1.19 (1.14-1.24)	3.51 (3.43-3.60)	1.20 (1.15-1.25)	0.11 (0.09-0.12)
40-64 yr (95% CI) ^{b)}	4.04 (3.95-4.13)	5.88 (5.77-5.99)	11.38 (11.22-11.53)	0.46 (0.43-0.49)
≥ 65 yr (95% CI) ^{b)}	4.99 (4.81-5.18)	7.11 (6.89-7.33)	28.68 (28.24-29.12)	0.94 (0.86-1.02)

CI, confidence interval; CNS, central nervous system; WHO, World Health Organization. ^{a)}Age-standardized incidence rate age-adjusted to the U.S. standard population in 2000, ^{b)}Age-specific incidence rate.