

Note to readers with disabilities: *EHP* strives to ensure that all journal content is accessible to all readers. However, some figures and Supplemental Material published in *EHP* articles may not conform to [508 standards](#) due to the complexity of the information being presented. If you need assistance accessing journal content, please contact ehp508@niehs.nih.gov. Our staff will work with you to assess and meet your accessibility needs within 3 working days.

Supplemental Material

Occurrence of Lead and Other Toxic Metals Derived from Drinking-Water Systems in Three West African Countries

Michael B. Fisher, Amy Z. Guo, J. Wren Tracy, Sridevi K. Prasad, Ryan D. Cronk, Emily G. Browning, Kaida R. Liang, Emma R. Kelly, and Jamie K. Bartram

Table of Contents

Table S1. Positive predictive value (PPV), negative predictive value (NPV), and percent agreement between test and retest WHO GV exceedance results for TMs in test-retest comparisons of water samples from the same water systems (N=109).

Table S2. Mean, median, and maximum relative mass fraction of 13 metals from water system component scrapings (N=61 systems).

Table S3. Multivariable linear regressions for associations between concentrations of TM of interest and other measured metals (controlling for country, pH, conductivity, and stagnation time).

Table S4. Normalized multivariable linear regression for associations between concentrations of lead and other measured metals (controlling for country, pH, conductivity, and stagnation time).

Table S5. Multivariable regression of log lead occurrence controlling for log lead groundwater occurrence and controlling for other relevant covariates (pH, conductivity, water system type, water system age, stagnation time, and country).

Table S6. Comparison of Ghana and Mali Samples to UNC Reanalysis of Duplicate Aliquots.

Figure S1. Indicative Calibration Curves.

Additional File- Excel Documents