

S3 Table. Procedural and outcome characteristics

Preoperative Iv Thrombolysis n (%)		31 (59.6)
Form of sedation/anesthesia n (%)	Conscious sedation	39 (75.0)
	General anesthesia	5 (9.6)
	Topical anesthesia	8 (15.4)
Safe onset to reperfusion (min), median (IQR), (n=28)		290 (206-354)
Safe onset to imaging (min), median (IQR), (n=35)		159(91-242)
Imaging to reperfusion time (min) median (IQR) (n=80) *		113 (90-155)
	Under median time n (%)	21 (40.4)
	In median time or over or no reperfusion n (%)	31 (59.6)
TICI Score	TICI 2b/3 n (%)	34 (65.4)
	TICI 0-2a, n (%)	18 (34.6)
Intra-arterial procedure	Retriever n (%)	35 (67.3)
	Aspiration only n (%)	11 (21.2)
	Clot not reached n (%)	6 (11.5)
Follow-up CT delay (days), mean \pm SD		1 \pm 0.75
FIV seen on Follow-up CT (mL) median (IQR)		26.1 (6.4-82.2)
Infarct Size over 70 mL n (%)		14 (26.9)

*11 Patients with TICI 0 excluded. Abbreviations: IQR=Interquartile Range, CTP=Computed Tomography Perfusion, TICI=Thrombolysis in Cerebral Infarction Scale, SD=Standard deviation, CT=Computed Tomography. IC=Ischemic Core, FIV=Final Infarct Volume