


C subunit of the ATP synthase is an amyloidogenic calcium dependent channel-forming peptide with possible implications in mitochondrial permeability transition

Giuseppe Federico Amodeo¹, Brenda Yasia Lee², Natalya Krilyuk¹, Carina Teresa Filice^{2,3}, Denis Valyuk¹, Daniel Erik Otzen⁴, Sergey Noskov⁵, Zoya Leonenko^{2,3}, Evgeny V. Pavlov^{1,*}


¹Department of Molecular Pathobiology, New York University,
²Department of Physics & Astronomy, University of Waterloo, Canada, ³Department of Biology, University of Waterloo, Canada, ⁴Interdisciplinary Nanoscience Center (iNANO) and Department of Molecular Biology and Genetics, Aarhus University, Aarhus C, Denmark, ⁵Centre for Molecular Simulation and Biochemistry Research Cluster, Department of Biological Sciences, University of Calgary, Canada

A


Exp. Time 40 sec

B


Exp. Time 10 sec

C


Exp. Time 20 sec

D

Exp. Time 30 sec

E

Exp. Time 50 sec

F

Exp. Time 60 sec

Supplementary Figure 1: Dot blots of the unmodified c subunit incubated in absence and in presence of Ca^{2+} using (A) anti c subunit antibodies and (B-F) A11-19 antibodies at different exposures (reported below every blot).