

Supplementary Table 2. HLA-DR expression in clones, obtained from freshly isolated unsorted NK cells and preliminary sorted HLA-DR-negative NK cell subset, on the 3rd week of their growth. HLA-DR surface expression MFI (normalized) was calculated as $(\text{exp}/\text{control} - 1) * 100$, where exp – mean of fluorescence intensity of experimental sample, control – mean of autofluorescence.

Clones from unsorted NK cells			Clones from HLA-DR-negative subset		
Clone #	HLA-DR-positive cells, %	HLA-DR-positive cells, MFI	Clone #	HLA-DR-positive cells, %	HLA-DR-positive cells, MFI
1g6	93.7	960.0	1g2	100.0	1868.8
2f5	96.6	993.3	1f7	100.0	2009.4
2b4	93.9	833.3	3g2	100.0	6868.8
2g9	94.7	1200.0	2g5	100.0	275.0
1g5	99.2	2896.7	5d4	100.0	7087.5
1f8	87.8	673.3	6e5	47.0	681.3
2b7	96.4	1100.0	3f6	61.0	587.5
1f10	90.5	683.3	4c5	37.0	618.8
2d7	99.6	1636.7	7d2	85.0	1150.0
2e3	76.4	430.0	8c4	100.0	962.5
2d2	99.8	1840.0	1-1	100.0	587.5
2c9	89.9	3123.3	1-2	33.0	1275.0
1e8	97.5	1090.0	1-3	23.0	275.0
1f11	64.8	316.7	1-4	100.0	4525.0
2e11	99.9	2970.0	1-5	100.0	743.8
1e11	100.0	5886.7	1-6	43.0	462.5
2g6	96.0	1266.7	1-7	100.0	2612.5
1b8	97.3	1450.0	1-8	100.0	306.3
1e6	99.6	1443.3	2-1	20.0	1587.5
1g4	36.0	140.0	2-2	20.0	1462.5
2g4	69.0	346.7	2-3	24.0	1556.3
			2-4	20.0	1775.0
			2-5	12.0	1587.5
			2-6	86.0	1993.8