

SUPPLEMENTAL MATERIAL

Table I: Summary of literature describing the incidence of stroke amongst COVID-19 patients.

Reference	Country	Types of Stroke	Incidence/Age
Yaghi et al ¹⁵	USA	Large and small vessel occlusion	0.9%/ Median=62.5 years
Mao et al ⁶	China	Ischemic stroke and cerebral haemorrhage	5.7%/ Mean= 52.7 years
Jain et al ⁸⁶	USA	Large/small vessel occlusion, ICH and hypoxic	1.1%/ Median=66 years
Klok et al ¹⁴	Netherlands	Ischaemic stroke (site not defined)	2.7%/ Mean= 64 years
Khandelwal et al ⁸⁷	USA, Spain and Italy	Large vessel occlusion	1.3%/ Mean =51 years
Bach et al ¹²	USA	Large-vessel occlusion and high-grade large-vessel stenosis	2.9%/ Mean = 63 years
Li et al ⁸⁸	China	Large/small vessel occlusion and ICH	4.6%/ 57-91 years
Siegler et al ⁸⁹	USA, Spain, Egypt, Romania	Large/small vessel occlusion and ICH	1.13%/ 60-79 years
Richter et al ⁹⁰	Germany	Not defined	1.2%/ Mean= 76.1 years
Lodigiani et al ⁹¹	Italy	Ischaemic stroke (site not defined)	2.5%/ Median =66 years
Fridman et al ⁵	Canada, USA, and Iran	Large/small vessel occlusion	1.8%/ N.A
Kvernland et al ⁹²	USA	ICH	0.5%/ Median= 60 years
Merkler et al ⁹³	USA	Ischaemic stroke (site not defined)	1.6%/ Median =69 years
Modin et al ⁹⁴	Denmark	Ischaemic stroke (site not defined)	0.86%/ Mean= 77 years