

Extracellular Vesicles Derived from Ascitic Fluid Enhance Growth and Migration of Ovarian Cancer Cells

Aparna Mitra¹, Kyoko Court¹, Travis Solley¹, Megan Mikkelsen¹, Chi Lam Au Yeung², Alpa Nick³, Karen Lu¹ and Ann H. Klopp^{1*}


1 Department of Radiation Oncology, The University of Texas MD Anderson Cancer Center, 1515 Holcombe Boulevard, Houston, TX 77030, USA.

Electronic address: aklopp@mdanderson.org.


2Department of Gynecologic Oncology and Reproductive Medicine, The University of Texas MD Anderson Cancer Center, Houston, TX 77030, USA.

3 Tennessee Oncology, Nashville, Tennessee 37203, USA.


Supplemental I:




Malignant Ascites S35




Malignant Ascites S35 + Benign exosome (S10)




Malignant Ascites S35 + Malignant exosome (S16)



Malignant Ascites S16




Malignant Ascites S16 + Malignant exosome (S35)




Malignant Ascites S16 + Malignant exosome (S16)

Supplemental II:



No peaks detected at 28S and 18S positions of the exosomal miRNAs





BAE
(S10) BAE
(S45) MAE
(S16) MAE
(S35)




Supplemental III:









Anti-CD 63, Non reducing




Anti-CD 9, Non reducing



Anti-CD 81, Non reducing



Anti-Alix , high exposure



Anti-Alix , low exposure