

OMTN, Volume 24

Supplemental information

**m⁶A-related lncRNAs are potential
biomarkers for predicting prognoses and
immune responses in patients with LUAD**

Feng Xu, Xiaoling Huang, Yangyi Li, Yongsong Chen, and Ling Lin

Figure S1




Figure S1. (A-C) Kaplan-Meier survival curves of the relative DSS, PFI, and DFI between the high- and low-risk groups.

Figure S2




Figure S2. (A-J) Identification of novel candidate compounds targeting the m6A-related lncRNA model

Table S1. Summary of patients' characteristics

Characteristics	Entire set		Training set		Testing set		p-Value
	Number	Percentage	Number	Percentage	Number	Percentage	
Age							
<60 year	136	26.98	89	29.28%	47	23.5	0.2025
≥60 year	358	71.03	210	69.08	148	74	
Not available	10	1.98	5	1.64	5	2.5	
Gender							
Female	270	53.57	165	54.28	105	52.5	0.7642
Male	234	46.43	139	45.72	95	47.5	
Stage							
Stage I-II	389	77.18	236	77.63	153	76.5	0.8231
Stage III-IV	107	21.23	63	20.72	44	22	
Not available	8	1.59	5	1.64	3	1.5	
T							
T1-2	437	86.71	269	88.49	168	84	0.1648
T3-4	64	12.7	33	10.86	31	15.5	
Not available	3	0.6	2	0.66	1	0.5	
M							
M0	335	66.47	203	66.78	132	66	0.8093
M1	25	4.96	14	4.61	11	5.5	
Not available	144	28.57	87	28.62	57	28.5	
N							
N0	325	64.48	197	64.8	128	64	0.9517
N1-3	167	33.13	100	32.89	67	33.5	
Not available	12	2.38	7	2.3	5	2.5	