

## Supplementary Material

**Table S1. IUD Counseling and IUD Method Choice by Participant Age, History of *N gonorrhoeae* or *C trachomatis* and Multiple Partners (Odds Ratios from Adjusted GEE models)**

	Counseled on IUD (n=1335)			Chose IUD Method (n=1342)		
	%	aOR (95% CI)	P	%	aOR (95% CI)	P
<b>Age Group</b>						
Adolescent	41.6	0.74 (0.53-1.02)	0.065	11.4	0.66 (0.46-0.95)	0.025
Young adult	51.9	REF	-	17.9	REF	-
<b>History of GC/CT</b>						
Yes	51.7	1.10 (0.83-1.47)	0.83	21.1	1.15 (0.81-1.63)	0.426
No	49.7	REF	-	15.5	REF	-
<b>Multiple partners, past 3m</b>						
Yes	46.3	0.80 (0.61-1.05)	0.101	12.4	0.66 (0.46-0.95)	0.025
No	50.5	REF	-	17.4	REF	-
<b>Study arm</b>						
Intervention	62.4	3.29 (2.37-4.56)	<0.001	19.0	1.79 (1.28-2.50)	0.001
Control	34.4	REF	-	13.3	REF	-

*Covariates included: race/ethnicity, insurance status, parity, prior LARC usage, practice setting. GC/CT: N gonorrhoeae or C trachomatis. CI: Confidence Intervals.*