

On-line Table 1. Distribution of examinations

	Newborns (N = 34)	Infants (N = 10)	Older Children (N = 27)
CT/CTV	19	4	7
MRI/MRV	12	6	17
CT/CTV & MRI/MRV	3	0	3

Note:—CTV indicates CT venography; MRI/MRV, MR imaging/MR venography.

On-line Table 3. Location of brain lesions

	Newborns (N = 34)	Infants (N = 10)	Older Children (N = 27)
Frontal lobe	9	2	6
Parietal lobe	8	1	2
Temporal lobe	5	1	0
Occipital lobe	4	0	1
Basal ganglia	3	0	0
Thalamus	3	0	3
Cerebellar hemisphere	0	2	0

On-line Table 2. Location of cerebral venous thrombosis in children

	Newborns (N = 34)	Infants (N = 10)	Older Children (N = 27)
Transverse sinus	25	9	14
Superior sagittal sinus	14	7	15
Straight sinus	12	3	4
Torcular	10	0	0
Sigmoid sinus	4	2	8
Internal cerebral vein	5	0	2
Vein of Galen	4	0	3
Medullary vein	3	0	0
Internal jugular vein	0	0	4

On-line Table 4. Size of brain lesions

	Newborns (N = 34)	Infants (N = 10)	Older children (N = 27)
<1 cm (small)	7 (33%)	1 (25%)	2 (17%)
1–3 cm (medium)	12 (57%)	1 (25%)	4 (33%)
>3 cm (large)	2 (10%)	2 (25%)	6 (59%)
Total	21	4	12