

Journal: *Drugs - Real World Outcomes*

Article title: Real-World Utilization and Safety of Daratumumab IV Rapid Infusions Administered in a Community Setting: A Retrospective Observational Study

Running head: Real-world utilization of daratumumab rapid infusions

Authors: Lucio Gordan, MD¹; Melody Chang, RPh¹; Marie-Hélène Lafeuille, MA²; Hela Romdhani, PhD²; Fuad Paramasivam, BA²; Eric M. Maiese, PhD³, Caroline McKay, PhD³

¹ Florida Cancer Specialists, Gainesville Florida, United States

² Analysis Group, Inc., Montreal Quebec, Canada

³ Janssen Scientific Affairs, LLC, Horsham New Jersey, United States

Corresponding author email: Marie-Helene.Lafeuille@analysisgroup.com

Supplemental Table 1. Non-infusion reaction adverse events^a

	N = 147
Number of patients with ≥ 1 non-IR AE, n (%)	15 (10.2%)
Number of non-IR AEs per patient,^b mean \pm SD [median; IQR]	1.9 \pm 1.2 [2;2]
Non-IR AEs, n (%)	
Gastrointestinal toxicities	4 (2.7%)
Constipation	1 (0.7%)
Diarrhea	1 (0.7%)
Nausea	2 (1.4%)
Vomiting	2 (1.4%)
General disorders and administration site conditions	8 (5.4%)
Fatigue	8 (5.4%)
Pyrexia	1 (0.7%)
Edema peripheral	0 (0.0%)
Chills	0 (0.0%)
Laboratory abnormalities	1 (0.7%)
Anemia	0 (0.0%)
Cytopenia	0 (0.0%)
Lymphopenia	0 (0.0%)
Neutropenia	1 (0.7%)
Thrombocytopenia	0 (0.0%)
Metabolism and nutrition disorder	1 (0.7%)
Decreased appetite	1 (0.7%)

Musculoskeletal and connective tissue disorders	2 (1.4%)
Muscle spasms	1 (0.7%)
Back pain	0 (0.0%)
Arthralgia	1 (0.7%)
Musculoskeletal chest pain	0 (0.0%)
Nervous system disorder	0 (0.0%)
Headache	0 (0.0%)
Peripheral sensory neuropathy	0 (0.0%)
Respiratory, thoracic and mediastinal disorders	0 (0.0%)
Cough	0 (0.0%)
Dyspnea	0 (0.0%)
Nasal congestion	0 (0.0%)
Vascular disorder	0 (0.0%)
Hypertension	0 (0.0%)
Hypotension	0 (0.0%)
Thrombosis	0 (0.0%)
Congestive heart failure	0 (0.0%)
Renal toxicities	0 (0.0%)
Other ^c	6 (4.1%)

Abbreviations: AE: adverse event; IQR: interquartile range; IR: infusion reaction; SD: standard deviation.

Notes:

- a. AEs were defined as health events explicitly attributed to daratumumab in the patient charts that did not also meet the definition of an infusion reaction.
- b. Among patients with at least one AE.
- c. The following other symptoms were reported: nosebleeds, GI bleeding, shakiness, dizziness, aches and pain, and abdominal pain.