http://dx.doi.org/10.1590/1516-4446-2020-0864

Stigma toward individuals with mental disorders among Brazilian psychiatrists: a latent class analysis - da Silva AG et al.

Table S1 Comparison of sociodemographic, professional, and clinical variables between the two highest stigma groups for schizophrenia and ADHD

Variable	Highest stigma - ADHD (n=174)	Highest stigma - schizophrenia (n=202)	p-value
Sociodemographics	, ,		•
Sex			
Female	75 (43.1)	88 (43.8)	0.89
Male	99 (56.9)	113 (56.2)	
Age, median (IQI)	40 (34/52)	41 (34/54)	0.48*
Marital status	,	,	
Married	114 (65.9)	140 (69.3)	0.48
Not married	59 (34.1) [′]	62 (30.7)	
Religion	(-)	- ()	
Yes	130 (74.7)	143 (70.8)	0.39
No	44 (25.3)	59 (29.2)	0.00
Professional			
Years since graduation, median (IQI)	14 (7/27)	15 (8/30)	0.31*
Graduate level education	11(1/21)	10 (0/00)	0.01
Yes (MSc and/or PhD)	36 (20.7)	56 (27.7)	0.11
No	138 (79.3)	146 (72.3)	0.11
Work, predominantly	100 (70.0)	140 (72.0)	
Private	118 (67.8)	148 (73.3)	0.25
Public	56 (32.2)	54 (26.7)	0.20
Teaching activities [†]	30 (32.2)	34 (20.7)	
Yes	96 (55.5)	115 (57.2)	0.74
No	77 (44.5)	86 (42.8)	0.74
NO	77 (44.5)	00 (42.0)	
Clinical		(()	
STAI-Trait total scores, median (IQI)	34 (29/40)	35 (29/40)	0.89*
PANAS positive, mean (SD)	34.4 (6.2)	33.8 (6.2)	0.29
PANAS negative, median (IQI)	17 (14/20)	16 (14/21)	0.49*
Big Five Inventory, mean (SD)			
Openness	3.7 (3.1/4.2)	3.6 (0.6)	0.44
Conscientiousness	3.7 (0.6)	3.6 (0.6)	0.29
Extroversion	3.2 (0.7)	3.3 (0.8)	0.60
Agreeableness	3.8 (0.5)	3.8 (0.5)	0.43
Neuroticism	2.5 (0.7	2.4 (0.7)	0.87
Family history of psychiatric treatment	•		
Yes	117 (68)	136 (67.7)	0.94
No	55 (32)	65 (32.3)	

Data presented as n (%), unless otherwise specified.

IQI = interquartile interval; PANAS = Positive and Negative Affect Schedule; SD = standard deviation; STAI = State-Trait Anxiety Inventory - Trait subscale.

^{*} Mann-Whitney *U* test.

[†] Undergraduate or graduate.