

Note to readers with disabilities: *EHP* strives to ensure that all journal content is accessible to all readers. However, some figures and Supplemental Material published in *EHP* articles may not conform to [508 standards](#) due to the complexity of the information being presented. If you need assistance accessing journal content, please contact ehp508@niehs.nih.gov. Our staff will work with you to assess and meet your accessibility needs within 3 working days.

Supplemental Material

Sex-Specific Effects of Prenatal and Early Life Inorganic and Methylated Arsenic Exposure on Atherosclerotic Plaque Development and Composition in Adult ApoE^{-/-} Mice

Luis Fernando Negro Silva, Kiran Makhani, Maryse Lemaire, Catherine A. Lemarié, Dany Plourde, Alicia M. Bolt, Christopher Chiavatti, D. Scott Bohle, Stéphanie Lehoux, Mark S. Goldberg, and Koren K. Mann

Table of Contents

Table S1. Effect of arsenicals on plasma cholesterol, triglycerides, HDL, LDL, ALT and AST levels of apoE^{-/-} male mice.

Table S2. Effect of arsenicals on plasma cholesterol, triglycerides, HDL, LDL, ALT and AST levels of apoE^{-/-} female mice.

Table S3. Comparison between the control groups of apoE^{-/-} and apoE^{-/-} As3mt^{-/-} DKO mice.

Table S4. Mean and standard deviations for the apoE^{-/-} dataset pertaining to the lesion areas in aortic arch and the aortic sinus.

Table S5. Mean and standard deviations for the apoE^{-/-} dataset pertaining to the macrophage, lipid, smooth muscle cell and collagen content in the aortic sinus.

Table S6. Mean and standard deviations for the apoE^{-/-} As3mt^{-/-} dataset pertaining to the lesion areas in aortic arch and the aortic sinus.