

Applied Microbiology and Biotechnology

Supplemental Information file

Process Development and Scale-up Optimization of the SARS-CoV-2 Receptor Binding Domain-Based Vaccine Candidate, RBD219-N1C1

Jungsoon Lee^{a,c*}, Zhuyun Liu^{a,c}, Wen-Hsiang Chen^{a,c}, Junfei Wei^{a,c}, Rakhi Kundu^{a,c},
Rakesh Adhikari^{a,c}, Joanne Altieri Rivera^{a,c}, Portia M. Gillespie^{a,c}, Ulrich Strych^{a,c},
Bin Zhan^{a,c}, Peter J. Hotez^{a, b,c,d,e}, Maria Elena Bottazzi^{a, b,c,d*}

^a *National School of Tropical Medicine, Department of Pediatrics, Baylor College of Medicine, One Baylor Plaza, BCM113 Houston, TX, 77030, USA*

^b *National School of Tropical Medicine, Department of Molecular Virology & Microbiology, Baylor College of Medicine, One Baylor Plaza, BCM113 Houston, TX, 77030, USA*

^c *Texas Children's Hospital Center for Vaccine Development, Baylor College of Medicine, 1102 Bates Street, Houston TX, 77030, USA*

^d *Department of Biology, College of Arts and Sciences, Baylor University, Waco TX, USA*

^e *James A. Baker III Institute for Public Policy, Rice University, Houston TX, USA*

*Correspondence to Maria Elena Bottazzi (Email: bottazzi@bcm.edu) Tel. 832-824-0504 Fax 832-825-0549 and Jungsoon Lee (Email: jslee@bcm.edu) Tel. 832-824-0504 Fax 832-825-0549

Supplementary Fig. S1




Fig. S1. Dynamic light scattering results of the purified proteins from Process-1 (A), Process-2 (B), and Process-3 (C). Measured Stokes radii, polydispersity (PD), and molecular weights (Mw) are shown as an average \pm SD from four independent measurements.