


Study	Sample size	Proportion (%)	95% CI	
Stelloo et al. 2016	219	8.676	5.304 to 13.217	Stelloo et al. 2016
An et al. 2007	17	41.176	18.444 to 67.075	An et al. 2007
Arabi et al. 2009	25	56.000	34.928 to 75.598	Arabi et al. 2009
Black et al. 2006	93	32.258	22.932 to 42.749	Black et al. 2006
Nagle et al. 2018	163	33.742	26.534 to 41.554	Nagle et al. 2018
Kommos et al. 2018	125	20.000	13.381 to 28.092	Kommos et al. 2018
Ruiz et al. 2014	64	9.375	3.519 to 19.297	Ruiz et al. 2014
Shikama et al. 2016	62	43.548	30.991 to 56.743	Shikama et al. 2016
Talhok et al. 2016	12	58.333	27.667 to 84.835	Talhok et al. 2016
Talhok et al. 2017	62	45.161	32.481 to 58.320	Talhok et al. 2017
Total (fixed effects)	842	24.654	21.793 to 27.691	Total (fixed effects)
Total (random effects)	842	32.105	21.371 to 43.896	Total (random effects)
Q	104.399			
DF	9			
Significance level	P < 0.001			
I ² (inconsistency)	91.38%			
95% CI for I ²	86.270 to 94.590			



Supplementary Fig. S11. Lymphovascular invasion analysis in mismatch repair deficiency in endometrial carcinoma [2,3,4,13,15-21,22,24,25]. CI, confidence interval.