


High FIB4 index is an independent risk factor of diabetic kidney disease in type 2 diabetes

Haruka Saito¹, Hayato Tanabe¹, Akihiro Kudo¹, Noritaka Machii¹, Moritake Higa², Satoshi Yamaguchi^{1,5}, Gulinu Maimaituxun¹, Kazumichi Abe³, Atsushi Takahashi³, Kenichi Tanaka⁴, Koichi Asahi⁶, Hiroaki Masuzaki⁷, Hiromasa Ohira³, Junichiro J. Kazama⁴ and Michio Shimabukuro^{1*}

¹Department of Diabetes, Endocrinology and Metabolism, ³Department of Gastroenterology and ⁴Department of Nephrology and Hypertension, Fukushima Medical University, Fukushima, Japan; ²Department of Diabetes and Lifestyle-Related Disease Center, Tomishiro Central Hospital, Okinawa, Japan; ⁵Department of Cardiology, Nakagami Hospital, Okinawa, Japan; ⁶Division of Nephrology and Hypertension, Iwate Medical University, Iwate, Japan; ⁷Division of Endocrinology, Diabetes and Metabolism, Hematology, Rheumatology (Second Department of Internal Medicine), University of the Ryukyus, Okinawa, Japan.

*Corresponding author at: Department of Diabetes, Endocrinology and Metabolism, Fukushima Medical University, 1 Hikarigaoka, Fukushima City, Fukushima 960-1295, Japan; E-mail address: mshimabukuro-ur@umin.ac.jp (M. Shimabukuro); Tel +81-24-547-1305 , Fax +81-24-547-1311



Supplement Figure 4. Pattern of missing data. In the left panel, green and yellow represents missing and existing values in the dataset. Each row represents the missing pattern. In the right panel, the horizontal bars display the frequencies of the corresponding combinations of existing and missing variables. VIM package 5.1.1 and ggplot 2 3.3 running on R 3.6.3. were used for visualization of the missing pattern.