


High FIB4 index is an independent risk factor of diabetic kidney disease in type 2 diabetes

Haruka Saito¹, Hayato Tanabe¹, Akihiro Kudo¹, Noritaka Machii¹, Moritake Higa², Satoshi Yamaguchi^{1,5}, Gulinu Maimaituxun¹, Kazumichi Abe³, Atsushi Takahashi³, Kenichi Tanaka⁴, Koichi Asahi⁶, Hiroaki Masuzaki⁷, Hiromasa Ohira³, Junichiro J. Kazama⁴ and Michio Shimabukuro^{1*}

¹Department of Diabetes, Endocrinology and Metabolism, ³Department of Gastroenterology and ⁴Department of Nephrology and Hypertension, Fukushima Medical University, Fukushima, Japan; ²Department of Diabetes and Lifestyle-Related Disease Center, Tomishiro Central Hospital, Okinawa, Japan; ⁵Department of Cardiology, Nakagami Hospital, Okinawa, Japan; ⁶Division of Nephrology and Hypertension, Iwate Medical University, Iwate, Japan; ⁷Division of Endocrinology, Diabetes and Metabolism, Hematology, Rheumatology (Second Department of Internal Medicine), University of the Ryukyus, Okinawa, Japan.

*Corresponding author at: Department of Diabetes, Endocrinology and Metabolism, Fukushima Medical University, 1 Hikarigaoka, Fukushima City, Fukushima 960-1295, Japan; E-mail address: mshimabukuro-ur@umin.ac.jp (M. Shimabukuro); Tel +81-24-547-1305 , Fax +81-24-547-1311



		Baseline	1	2	3	4	5	6	7	8	9	10	11
FIB4 >1.30	Median	1.84	1.86	1.88	1.96	2.02	2.06	2.14	2.13	2.06	2.18	2.11	2.24
	25%	1.57	1.48	1.55	1.62	1.61	1.65	1.64	1.72	1.67	1.69	1.62	1.79
	75%	2.32	2.49	2.51	2.53	2.53	2.60	2.73	2.60	2.80	2.73	2.77	2.90
FIB4 ≤1.30	Median	0.77	0.87	0.90	0.98	1.07	1.07	1.10	1.07	1.03	1.04	1.13	1.13
	25%	0.60	0.65	0.69	0.76	0.75	0.77	0.80	0.78	0.82	0.78	0.83	0.84
	75%	1.02	1.20	1.23	1.30	1.38	1.41	1.41	1.48	1.41	1.50	1.48	1.69

Supplement Figure 6. Annual changes in FIB4 index during the observational periods in FIB4 index >1.30 or ≤1.30. Values are median [25%, 75%].