


Supplementary material online**Supplementary Figure S1**

Distribution of propensity scores between statin and non-statin groups in the overall cohort.



Supplementary Table S1

Patient characteristics of the propensity score matched and unmatched groups

	Unmatched		P-value
	Matched group	group	
	n=1872	n=716	
Preprocedural variables			
Age, years	84.2±5.1	84.6±5.4	0.08
Male, n (%)	532 (28.4%)	263 (36.7%)	<0.01
Body surface area, m ²	1.43±0.17	1.42±0.17	0.16
NYHA class 3 or 4, n (%)	919 (49.1%)	402 (56.2%)	<0.01
Clinical Frailty Scale	3.8±1.2	4.2±1.4	<0.01
Smoking, n (%)	364 (19.4%)	160 (22.4%)	0.10
Diabetes mellitus, n (%)	402 (21.5%)	153 (21.4%)	0.95
Hypertension, n (%)	1497 (80.0%)	493 (68.9%)	<0.01
Chronic kidney disease, n (%)	1291 (69.0%)	518 (72.4%)	0.09
Atrial fibrillation, n (%)	364 (19.4%)	185 (25.8%)	<0.01

Coronary artery disease, <i>n</i> (%)	750 (40.1%)	204 (28.5%)	<0.01
Previous myocardial infarction, <i>n</i> (%)	96 (5.1%)	62 (8.7%)	<0.01
Previous percutaneous coronary intervention, <i>n</i> (%)	487 (26.0%)	139 (19.4%)	<0.01
Previous coronary artery bypass grafting, <i>n</i> (%)	117 (6.3%)	52 (7.3%)	0.36
Peripheral artery disease, <i>n</i> (%)	264 (14.1%)	113 (15.8%)	0.28
Previous stroke, <i>n</i> (%)	9 (0.5%)	3 (0.4%)	0.98
STS risk score	7.7±5.7	9.8±9.2	<0.01
Renin-angiotensin inhibitor, <i>n</i> (%)	1084 (57.9%)	303 (42.3%)	<0.01
Beta blocker, <i>n</i> (%)	637 (34.0%)	243 (33.9%)	0.97
eGFR, mL/min/1.73 m ²	52.0±19.1	49.7±20.1	<0.01
Haemoglobin, g/dL	11.3±1.6	11.2±1.7	0.16
Procedural variables			
Transfemoral approach, <i>n</i> (%)	1554 (83.0%)	613 (85.6%)	0.11

Local anaesthesia, <i>n</i> (%)	1448 (77.4%)	530 (74.0%)	0.08
Contrast volume, mL	116.1±58.6	114.1±58.2	0.43
Fluoroscopy time, min	21.4±11.4	21.2±10.5	0.66
Procedure time, min	81.0±43.3	80.3±50.9	0.72

CT, computed tomography; eGFR, estimated glomerular filtration rate; NYHA, New York Heart Association; STS, Society of Thoracic Surgeons.

Supplementary Table S2

In-hospital outcomes of the propensity score-matched and -unmatched groups

	Matched group <i>n</i> =1872	Unmatched group <i>n</i> =716	<i>P</i>-value
All-cause mortality, <i>n</i> (%)	43 (2.3%)	27 (3.8%)	<0.01
Acute kidney injury, <i>n</i> (%)	201 (10.7%)	88 (12.3%)	0.27
Bleeding, <i>n</i> (%)	421 (22.5%)	199 (27.8%)	<0.01
Stroke, <i>n</i> (%)	43 (2.3%)	18 (2.5%)	0.75
Vascular complication, <i>n</i> (%)	168 (9.0%)	65 (9.1%)	0.93