

Table S1. Clinicopathological associations of pIGF1R expression in triple-negative breast cancer

Clinicopathological parameters	pIGF1R Negative N=19 (%)	pIGF1R Positive N=30 (%)	P value*
Age at diagnosis, years	52.3±7.1	55.7±12.9	
≤50	8(42.1)	4(13.3)	0.0392
≥51	11(57.9)	26(86.7)	
Grade			
High	16(84.2)	16(53.3)	0.0718
Moderate	3(15.8)	8(26.7)	
Low	0(0)	6(20.0)	
T stage			
T1	6(31.6)	8(26.7)	0.8114
T2	12(63.1)	19(63.3)	
T3	1(5.3)	3(10.0)	
N stage			
N0	13(68.4)	21(70.0)	>0.05
N1	6(31.6)	9(30.0)	
M stage			
M0	19(100)	29(96.7)	>0.05
M1	0(0)	1(3.3)	
Ki67			0.0338
<14%	0(0)	7(23.3)	
≥14%	19(100)	23(76.7)	
Recurrence			
No	15(78.9)	21(70.0)	0.7408
Yes	4(21.1)	9(30.0)	
Median survival-up, months	43	28	

All cases were diagnosed as invasive ductal carcinoma and classified according to the American Joint Committee on Cancer (AJCC) staging system. T: tumor; N: lymph node; M: metastasis.

*P values calculated by Chi-square testing; bold if statistically significant (P < 0.05)