

Supplementary Table 1. Characteristics of Survey Respondents vs Nonrespondents

	Respondents (n = 1020)	Nonrespondents (n = 1851)	P value
Age			.415
21–50 y	143 (14.0)	273 (14.8)	
51–60 y	334 (32.7)	638 (34.6)	
61–90 y	543 (53.2)	935 (50.7)	
Male	649 (63.6)	1293 (70.0)	.001
Race/ethnicity			.001
Non-Hispanic White	359 (35.2)	773 (42.0)	
Non-Hispanic Black	292 (28.6)	507 (27.5)	
Hispanic	341 (33.4)	506 (27.5)	
Other	28 (2.7)	55 (3.0)	
Etiology of liver disease			.216
Hepatitis C virus	603 (59.1)	1023 (55.3)	
Alcohol related	179 (17.5)	351 (19.0)	
Nonalcoholic steatohepatitis	136 (13.3)	281 (15.2)	
Hepatitis B virus	28 (2.7)	41 (2.2)	
Other	74 (7.3)	155 (8.4)	
Child-Turcotte-Pugh class A	754 (73.9)	1287 (69.5)	.021
CirCom score ^a			.729
0	352 (34.5)	618 (33.4)	
1	285 (27.9)	499 (27.0)	
2	173 (17.0)	326 (17.6)	
3	38 (3.7)	86 (4.7)	
4	172 (16.9)	322 (17.4)	
Receipt of hepatology care ^a	694 (68.0)	873 (47.2)	<.0001

NOTE. Values are n (%).

CirCom, cirrhosis comorbidity.

^aDuring year prior to randomization.**Supplemental Table 2.** Prevalence of Depression and Anxiety, Stratified by Child-Turcotte-Pugh Class

	Parkland	UTSW	MEDVAMC	P value
Child-Turcotte-Pugh class A cirrhosis				
Depression measured by PHQ-9 score				
No-minimal depression (0–4)	173 (39.1)	60 (47.62)	72 (38.92)	.09
Mild depression (5–9)	104 (23.5)	37 (29.37)	48 (25.95)	
Moderate depression (10–14)	96 (21.7)	22 (17.46)	37 (19.46)	
Moderately severe depression (15–19)	52 (11.7)	5 (3.97)	18 (9.73)	
Severe depression (20–25)	18 (4.1)	2 (1.59)	11 (5.95)	
Anxiety measured by STAI score				
Mild anxiety (<40)	245 (55.3)	76 (60.3)	128 (69.2)	.005
Moderate–severe anxiety (≥40)	198 (44.7)	50 (39.7)	57 (30.8)	
Child Pugh Turcotte class B–C cirrhosis				
Depression measured by PHQ-9 score				
No-minimal depression (0–4)	65 (35)	12 (41.4)	14 (27.5)	.80
Mild depression (5–9)	49 (26.3)	8 (27.6)	16 (31.4)	
Moderate depression (10–14)	33 (17.7)	6 (20.7)	10 (19.6)	
Moderately severe depression (15–19)	30 (16.1)	3 (10.3)	7 (13.7)	
Severe depression (20–25)	9 (4.8)	0 (0)	4 (7.8)	
Anxiety measured by STAI score				
Mild anxiety (<40)	90 (48.4)	19 (65.5)	28 (54.9)	.20
Moderate–severe anxiety (≥40)	96 (51.6)	10 (34.5)	23 (45.1)	

NOTE. Values are n (%).

MEDVAMC, Michael E. DeBakey VA Medical Center; PHQ-9, Patient Health Questionnaire-9; STAI, State–Trait Anxiety Inventory; UTSW, University of Texas Southwestern Medical Center.