Synthesis and characterization of thiolated gum ghatti as a novel excipient: Development of compression coated mucoadhesive tablets of domperidone

Vivek Puri^{1,2}, Ameya Sharma^{1,2}, Pradeep Kumar³, Inderbir Singh^{2*}, Kampanart Huanbutta^{4*}

¹Chitkara College of Pharmacy, Chitkara University, Punjab 140401, India

²Chitkara University School of Pharmacy, Chitkara University, Himachal Pradesh 174103, India

³Wits Advanced Drug Delivery Platform Research Unit, Department of Pharmacy and Pharmacology, School of Therapeutic Sciences, Faculty of Health Sciences, University of the Witwatersrand, Johannesburg 2193, South Africa

⁴Faculty of Pharmaceutical Sciences, Burapha University, 169, Saensook, Muang, Chonburi, 20131, Thailand

*Corresponding Author: Email: <u>inderbir.singh@chitkara.edu.in</u>; Email: <u>kampanart@go.buu.ac.th</u>




Figure S1. FTIR spectra of pure drug and tablet components. (A) Drug (Domperidone); (B) Avicel-112; (C) PVP K30; (D) Talc; (E) Magnesium stearate; (F) Drug with avicel-112; (G) Drug with PVP K30; (H) Drug with talc; (I) Drug with magnesium stearate




Figure S2. UV absorption maxima of domperidone




Figure S3. Standard plot of thioglycolic acid