

Table S3. Sensitivity analyses of the associations of three thyroid markers and the risk of GDM according to the grounds GDM is diagnosed.

Trimester	Group	<i>OR</i>₁ (95%<i>CI</i>)	<i>OR</i>₂ (95%<i>CI</i>)	<i>OR</i>₃ (95%<i>CI</i>)
1st trimester				
	High fasting glucose	1.76 (1.25, 2.47)	0.96 (0.89, 1.04)	1.10 (1.05, 1.16)
	High postprandial glucose	1.66 (1.18, 2.33)	1.03 (0.97, 1.10)	1.08 (1.02, 1.14)
	Both	1.38 (0.80, 2.35)	0.96 (0.86, 1.09)	1.05 (0.97, 1.15)
2nd trimester				
	High fasting glucose	2.46 (1.72, 3.54)	0.89 (0.79, 1.00)	1.10 (1.06, 1.14)
	High postprandial glucose	2.40 (1.64, 3.49)	0.87 (0.77, 0.99)	1.09 (1.05, 1.13)
	Both	2.57 (1.47, 4.51)	0.89 (0.74, 1.07)	1.10 (1.05, 1.17)

Abbreviations: OR, odds ratio; CI, confidence interval;
*OR*₁ is for T3; *OR*₂ is for T3/FT4; *OR*₃ is for TSH.