

SUPPLEMENTARY MATERIAL

Sensitivity of Field Isolates of *Eimeria acervulina* and *E. maxima* from Three Regions in Brazil to Eight Anticoccidial Drugs

TABLE. DESCRIPTION OF THE SUPPLEMENTARY MATERIAL

PAGE	Parameter	Challenge	AST
2	Body Weight Gain (BWG)	<i>E. acervulina</i>	2 and 3
3	Body Weight Gain (BWG)	<i>E. acervulina</i>	2 and 3
4	Body Weight Gain (BWG)	<i>E. maxima</i>	1 and 4
5	Body Weight Gain (BWG)	<i>E. maxima</i>	1 and 4
6	Feed Conversion Ratio (FCR)	<i>E. acervulina</i>	2 and 3
7	Feed Conversion Ratio (FCR)	<i>E. acervulina</i>	2 and 3
8	Feed Conversion Ratio (FCR)	<i>E. maxima</i>	1 and 4
9	Feed Conversion Ratio (FCR)	<i>E. maxima</i>	1 and 4
10	Lesion Score (LS)	<i>E. acervulina</i>	2 and 3
11	Lesion Score (LS)	<i>E. acervulina</i>	2 and 3
12	Lesion Score (LS)	<i>E. maxima</i>	1 and 4
13	Lesion Score (LS)	<i>E. maxima</i>	1 and 4
14	OAA (POAA)	<i>E. acervulina</i>	2 and 3
15	OAA (POAA)	<i>E. acervulina</i>	2 and 3
16	OAA (POAA)	<i>E. maxima</i>	1 and 4
17	OAA (POAA)	<i>E. maxima</i>	1 and 4































