## Short-Term Suicide Risk Following Psychiatric Hospital Discharge

## Supplementary Material

Supplemental Table 1: Cohort selection among inpatients who were 18 to 64 years of age

Supplemental Table 2: Background characteristics of adult Medicaid inpatients with a principal diagnosis of depression, bipolar disorder, schizophrenia, substance use, or no mental disorder

Supplemental Table 3: Suicide during the 90 days following hospital discharge of adult inpatients in the Medicaid program by presence of primary mental disorder diagnosis and age group

Supplemental Table 4: Short-term risk of suicide of adult inpatients in the Medicaid program with a first-listed mental disorder

Supplemental Table 5: Suicide rates of adult inpatients in the Medicaid program by number of outpatient health visits in the 180 days before hospital admission, during the first 90 days following hospital discharge

Appendix (Diagnostic Codes)

| Supplemental Table 1: Cohort selection among inpatients who were  | 18 to 64 years of age        |                         |
|---|------------------------------|-------------------------|
| Eligibility criterion   | Any Mental Disorders         | Non-Mental Disorders    |
|   | Cohort (N)                   | Cohort (N)*             |
| Initial sample  | 1,163,328                    | 1,518,113               |
| Eligible for Medicaid, 180 days prior to index admission  | 901,641                      | 1,128,441               |
| Inpatient length of stay, 1 to 30 days  | 890,451                      | 1,128,192               |
| Discharged home, not an institutional setting   | 770,643                      | 1,090,551               |
| Study cohorts   | 770,643                      | 1,090,551               |
| Results based on 45-state 2001-2007 Medicaid MAX files. *10% ra other than mental disorders (ICD-9: 290-319). | ndom sample of inpatients wi | th principal discharges |

| Supplemental Table 2: Background characteristics of adult Medicaid inpatients with a principal diagnosis of depression, bipolar disorder, |
|---|
| ashizanhrania subatanga uga ar na mantal digardar   |

|  |                          |                         |                              | Substance                |                          | Non-Mental         |
|--|--------------------------|-------------------------|------------------------------|--------------------------|--------------------------|--------------------|
| Characteristic                           | Depressive               | Bipolar                 | Oakiaaakaaaia                | Use                      | Other Mental             | Disorders          |
|  | Disorders<br>(N=193,620) | Disorder<br>(N=109,139) | Schizophrenia<br>(N=173,810) | Disorders<br>(N=185,064) | Disorders<br>(N=109,010) | (N=1,090,551)<br>% |
|  | (N=193,020)<br>%         | (N=109,139)<br>%        | (N-173,610)<br>%             | (N=185,064)<br>%         | (N=109,010)<br>%         | 70                 |
| Age, years                               | ,,                       | ,,                      | ,,                           |                          | ,,,                      |                    |
| 18-34                                    | 39.7                     | 41.0                    | 30.8                         | 26.9                     | 44.8                     | 64.9               |
| 35-44                                    | 28.7                     | 29.6                    | 30.0                         | 35.2                     | 24.6                     | 12.5               |
| 45-64                                    | 31.6                     | 29.4                    | 39.2                         | 37.8                     | 30.6                     | 22.5               |
| Sex                                      |                          |                         |                              |                          |                          |                    |
| Male                                     | 30.0                     | 33.2                    | 55.6                         | 57.8                     | 41.5                     | 14.6               |
| Female                                   | 70.0                     | 66.8                    | 44.4                         | 42.2                     | 58.5                     | 85.4               |
| Race/Ethnicity                           |                          |                         |                              |                          |                          |                    |
| White, non-Hispanic*                     | 67.9                     | 76.5                    | 56.6                         | 55.2                     | 67.1                     | 51.8               |
| Black, non-Hispanic                      | 22.0                     | 17.4                    | 36.2                         | 33.7                     | 24.8                     | 27.6               |
| Hispanic                                 | 10.0                     | 6.1                     | 7.3                          | 11.1                     | 8.1                      | 20.6               |
| Recent mental health treatment**         |                          |                         |                              |                          |                          |                    |
| Outpatient mental health diagnosis       |                          |                         |                              |                          |                          |                    |
| Depression                               | 44.4                     | 28.1                    | 16.2                         | 17.9                     | 24.3                     | 4.6                |
| Alcohol use disorder                     | 11.9                     | 11.7                    | 8.8                          | 30.9                     | 9.3                      | 2.2                |
| Drug use disorder                        | 6.4                      | 6.4                     | 4.2                          | 25.7                     | 4.9                      | 1.0                |
| Schizophrenia                            | 5.3                      | 12.4                    | 55.8                         | 4.5                      | 9.6                      | 0.9                |
| Bipolar disorder                         | 7.9                      | 44.0                    | 10.5                         | 5.5                      | 9.1                      | 1.0                |
| Anxiety disorder                         | 16.7                     | 13.4                    | 5.9                          | 9.7                      | 14.7                     | 2.2                |
| Other mental disorder                    | 19.7                     | 22.8                    | 22.0                         | 10.9                     | 31.5                     | 3.3                |
| Any deliberate self-harm                 | 1.9                      | 1.4                     | 0.5                          | 0.4                      | 1.2                      | 0.0                |
| Emergency mental health diagnosis        | 20.6                     | 24.0                    | 22.4                         | 18.4                     | 19.9                     | 1.1                |
| Inpatient, length of stay, days          |                          |                         |                              |                          |                          |                    |
| 1-4                                      | 40.3                     | 29.4                    | 18.8                         | 47.7                     | 49.9                     | 77.0               |
| 5-8                                      | 39.1                     | 40.0                    | 36.0                         | 36.4                     | 30.5                     | 16.9               |
| 9-30                                     | 20.6                     | 30.6                    | 45.3                         | 15.8                     | 19.6                     | 5.4                |
| No outpatient health care, past 180 days | 8.8                      | 8.6                     | 11.1                         | 12.3                     | 10.3                     | 6.7                |

<sup>\*</sup>Includes White, American Indian/Alaskan Native, Asian, Native Hawaiian/Other Pacific Islander, and more than one race \*\*During 180 day period prior to inpatient admission

| Group                | Number of<br>Discharges<br>N | Person-<br>years<br>at risk<br>Years | Suicides<br>N | Suicide Rate<br>(per 100,000<br>person-<br>years) | Unadjusted<br>Hazard Ratio<br>of Suicide (95%CI) | Adjusted*<br>Hazard Ratio<br>of Suicide (95%CI) |
|----------------------|------------------------------|--------------------------------------|---------------|---|--|---|
| Age, total           |                              |                                      |               |   |  |   |
| Any mental disorder  | 770643                       | 189573.7                             | 338           | 178.3   | 15.43 (10.68, 22.28)                             | 10.61 (7.27, 15.48)                             |
| Non-mental disorder  | 1090551                      | 268214.9                             | 31            | 11.6  | 1.00   | 1.00  |
| Age, 18-34 years     |                              |                                      |               |   |  |   |
| Any mental disorder  | 273726                       | 67428.2                              | 90            | 133.5   | 46.59 (18.93, 114.65)                            | 21.44 (8.38, 54.83)                             |
| Non-mental disorder  | 708081                       | 174540.6                             | 5             | 2.9   | 1.00   | 1.00  |
| Age, 35-44 years     |                              |                                      |               |   |  |   |
| Any mental disorder  | 232176                       | 57123.6                              | 109           | 190.8   | 7.13 (3.61, 14.06)                               | 5.71 (2.88, 11.34)                              |
| Non- mental disorder | 136703                       | 33605.1                              | 9             | 26.8  | 1.00   | 1.00  |
| Age, 45-64 years     |                              |                                      |               |   |  |   |
| Any mental disorder  | 264741                       | 65021.9                              | 139           | 213.8   | 7.56 (4.57, 12.51)                               | 6.86 (4.13, 11.39)                              |
| Non-mental disorder  | 245767                       | 60069.2                              | 17            | 28.3  | 1.00   | 1.00  |

| Supplemental Table 4: Short-term risk of suici       | de of adult inpatients in        | the Medicaid program v           | with a first-listed mental         | disorder               |
|--|----------------------------------|----------------------------------|------------------------------------|------------------------|
| Characteristic                                       | Total<br>Adjusted<br>HR (95% CI) | Males<br>Adjusted<br>HR (95% CI) | Females<br>Adjusted<br>HR (95% CI) | Interaction<br>P-value |
| Age, year  | (6676 6.)                        | 1.11 (00 / 0 0.)                 | (00 / 00 / 01 /                    |                        |
| 18-34  | 1.0                              | 1.0                              | 1.0                                | N/A                    |
| 35-44  | 1.58 (1.19, 2.10)                | 1.25 (0.87, 1.81)                | 2.28 (1.43, 3.62)                  | 0.05                   |
| 45-64  | 1.79 (1.36, 2.35)                | 1.59 (1.13, 2.23)                | 2.22 (1.40, 3.54)                  | 0.25                   |
| Sex  |                                  |                                  |                                    | N/A                    |
| Male   | 2.29 (1.82, 2.88)                | N/A                              | N/A                                |                        |
| Female   | 1.00                             | N/A                              | N/A                                |                        |
| Race/Ethnicity                                       |                                  |                                  |                                    |                        |
| White, non-Hispanic*                                 | 4.91 (3.22, 7.47)                | 4.43 (2.68, 7.32)                | 6.09 (2.83, 13.15)                 | 0.49                   |
| Hispanic   | 2.23 (1.16, 4.27)                | 1.86 (0.82, 4.19)                | 3.16 (1.05, 9.50)                  | 0.49                   |
| Black, non-Hispanic                                  | 1.00                             | 1.00                             | 1.00                               | N/A                    |
| Primary discharge diagnosis                          |                                  |                                  |                                    |                        |
| Depressive disorders                                 | 1.99 (1.41, 2.81)                | 2.14 (1.40, 3.30)                | 1.72 (0.96, 3.08)                  | 0.64                   |
| Bipolar  | 1.65 (1.11, 2.45)                | 1.72 (1.04, 2.86)                | 1.46 (0.76, 2.80)                  | 0.79                   |
| Schizophrenia  | 1.37 (0.94, 2.00)                | 1.43 (0.91, 2.25)                | 1.19 (0.58, 2.42)                  | 0.75                   |
| Substance use disorder                               | 1.0                              | 1.0                              | 1.0                                | N/A                    |
| Other mental disorder                                | 1.38 (0.91, 2.08)                | 1.35 (0.80, 2.28)                | 1.31 (0.66, 2.61)                  | 0.98                   |
| Number of outpatient mental health disorder groups** |                                  |                                  |                                    |                        |
| 0  | 1.0                              | 1.0                              | 1.0                                | N/A                    |
| 1  | 1.15 (0.82, 1.60)                | 1.17 (0.76, 1.79)                | 1.09 (0.64, 1.87)                  | 0.83                   |
| 2  | 1.48 (1.04, 2.10)                | 1.39 (0.88, 2.20)                | 1.57 (0.91, 2.71)                  | 0.76                   |
| 3  | 1.39 (0.89, 2.18)                | 1.55 (0.88, 2.73)                | 1.16 (0.56, 2.41)                  | 0.51                   |
| ≥4   | 1.92 (1.13, 3.25)                | 1.64 (0.80, 3.36)                | 2.31 (1.05, 5.09)                  | 0.54                   |
| Linear Trend Test: Chi-square (p-value)              | 3.59 (0.06)                      | 1.12 (0.29)                      | 2.74 (0.10)                        |                        |
| Inpatient, length of stay, days                      |                                  |                                  |                                    |                        |
| 1-4  | 1.13 (0.84, 1.52)                | 1.05 (0.72, 1.54)                | 1.25 (0.79, 1.99)                  | 0.49                   |
| 5-8  | 0.99 (0.74, 1.33)                | 1.09 (0.76, 1.56)                | 0.85 (0.52, 1.38)                  | 0.47                   |
| 9-30   | 1.0                              | 1.0                              | 1.0                                | N/A                    |
| No outpatient health care, past 180 days             | 1.67 (1.11, 2.52)                | 1.53 (0.92, 2.56)                | 2.02 (1.01, 4.03)                  | 0.56                   |

HR denotes hazard ratios, CI denotes confidence interval. Adjusted HRs are from a single model that also controls for geographic region.
\*Includes White, American Indian/Alaskan Native, Asian, Native Hawaiian/Other Pacific Islander, and more than one race
\*\*180 days prior to hospital admission

Supplemental Table 5: Suicide rates of adult inpatients in the Medicaid program by number of outpatient health visits in the 180 days before hospital admission, during the first 90 days following hospital discharge Suicide Rate Unadjusted Adjusted\* Number of Person-(per 100,000 Hazard Ratio Hazard Ratio Group Discharges years Suicides person-years) of Suicide of Suicide (95%CI) Ν Ν at risk (95%CI) Years Outpatient health care visits, past 180 days 0 153211 37691.6 45 119.4 1.11 (0.76, 1.62) 1.40 (0.94, 2.07) 1-5 286531 70513.9 50 70.9 0.66 (0.46, 0.95) 0.88 (0.61, 1.29) 6-10 237982 58584.4 38 64.9 0.60 (0.41, 0.90) 0.85 (0.56, 1.27) 11-15 217741 53603.6 35 65.3 0.61 (0.40, 0.92) 0.92 (0.61, 1.40) 16-20 44416.4 30 67.5 0.93 (0.60, 1.45) 180409 0.63 (0.41, 0.97) 42 21-30 263174 64778.7 64.8 0.60 (0.41, 0.89) 0.86 (0.58, 1.28) 31-50 267833 65884.3 62 94.1 0.88 (0.62, 1.24) 1.17 (0.82, 1.67) 67 107.5 >50 254313 62315.7 1.00 1.00 Adjusted for age, sex, race/ethnicity, and region.

| Appendix: Diagnostic Groups and ICD-9-CM Code | S   |
|---|---|
| Diagnostic Group and Conditions               | ICD-9-Codes   |
| Depressive disorders                          | 296.2x, 296.3x, 296.82, 298.0, 300.4, 311                     |
| Alcohol use disorders                         | 291, 303, 305   |
| Drug use disorders                            | 292, 304  |
| Schizophrenia                                 | 295.00-295.99   |
| Bipolar disorders                             | 296.0x, 296.1x, 296.4x, 296.5x, 296.6x, 296.7x, 296.8x (other |
|   | than 296.82)  |
| Anxiety Disorders                             | 300.0x, 300.2x, 300.3 308.3, 309.81                           |
| Other mental disorders                        | 290-319 not included in other narrow diagnostic groups        |
| Deliberate self-harm                          | E950-E959   |

Diagnostic groups define diagnostic categories used in Table 3 and Supplemental Table 2. Substance use disorders in all tables defined as combination of alcohol use disorders and drug use disorders.