


Anlotinib, Vincristine and Irinotecan (AVI) for Advanced Ewing Sarcoma after Failure of Standard Multimodal Therapy: A Two-cohort, Phase Ib/II Trial

Wei Guo et al.








Anlotinib, Vincristine and Irinotecan (AVI) for Advanced Ewing Sarcoma after Failure of Standard Multimodal Therapy: A Two-cohort, Phase Ib/II Trial

Wei Guo et al.



Change From Baseline in Target Lesion

Figure S1B.



Supplemental Figures for: Anlotinib, Vincristine and Irinotecan (AVI) for Advanced Ewing Sarcoma after Failure of Standard Multimodal Therapy: A Two-cohort, Phase Ib/II Trial Wei Guo et al.

Figure

Figure S2a.



Anlotinib, Vincristine and Irinotecan (AVI) for Advanced Ewing Sarcoma after Failure of Standard Multimodal Therapy: A Two-cohort, Phase Ib/II Trial

Wei Guo et al.




Figure S2b



Anlotinib, Vincristine and Irinotecan (AVI) for Advanced Ewing Sarcoma after Failure of Standard Multimodal Therapy: A Two-cohort, Phase Ib/II Trial

Wei Guo et al.




Figure S3.



Anlotinib, Vincristine and Irinotecan (AVI) for Advanced Ewing Sarcoma after Failure of Standard Multimodal Therapy: A Two-cohort, Phase Ib/II Trial

Wei Guo et al.




Figure S4.