

**Appendix 1. Sensitivity analyses predicting loss to follow-up (LTFU) by pre-disaster psychiatric disorder: The PREDICT study (2003-2011).**

<i>Pre-disaster Disorder</i>	<i>n among those LTFU (n=941)*</i>	<i>n among those not LTFU (n=1,708)**</i>	$\chi^2$	<i>p</i>
<b>Eating Disorders</b>				
Yes	8	17	0.137	0.712
No	933	1,691		
<b>Anorexia</b>				
Yes	1	0	1.817	0.178
No	939	1,707		
<b>Bulimia</b>				
Yes	7	17	0.427	0.513
No	934	1,691		
<b>Mood Disorders</b>				
Yes	260	520	2.314	0.128
No	681	1,188		
<b>Major Depressive</b>				
Yes	219	417	0.323	0.570
No	712	1,284		
<b>Bipolar I</b>				
Yes	16	36	0.523	0.469
No	925	1,672		
<b>Dysthymia</b>				

Yes	59	133	2.098	0.148
No	873	1,557		
<hr/>				
Psychotic Disorders				
Yes	27	45	0.130	0.718
No	910	1,658		
<hr/>				
Schizophrenia				
Yes	4	6	0.076	0.783
No	911	1,632		
<hr/>				
Schizophreniform				
Yes	1	1	0.184	0.668
No	931	1,692		
<hr/>				
Schizoaffective				
Yes	1	2	0.007	0.935
No	931	1,686		
<hr/>				
Delusional				
Yes	1	0	1.798	0.180
No	921	1,657		
<hr/>				
Brief Psychotic				
Yes	21	37	0.014	0.904
No	868	1,581		
<hr/>				
Anxiety Disorders (not including PTSD)				
Yes	332	707	9.507	0.002

No	609	1,001		
<hr/>				
Obsessive Compulsive				
Yes	19	24	1.367	0.242
No	885	1,603		
<hr/>				
Panic				
Yes	53	75	2.015	0.156
No	883	1,622		
<hr/>				
Agoraphobia				
Yes	47	70	1.096	0.295
No	885	1,613		
<hr/>				
Social Phobia				
Yes	47	90	0.092	0.761
No	881	1,595		
<hr/>				
Generalized Anxiety				
Yes	9	20	0.253	0.615
No	928	1,684		
<hr/>				
Specific Phobia				
Yes	300	659	11.658	0.001
No	637	1,044		
<hr/>				
Somatoform Disorders				
Yes	107	177	0.644	0.422
No	834	1,531		
<hr/>				
Somatization				

Yes	2	1	1.269	0.260
No	939	1,705		
<hr/>				
Conversion				
Yes	55	89	0.357	0.550
No	829	1,491		
<hr/>				
Pain				
Yes	64	104	0.505	0.477
No	803	1,467		
<hr/>				
Hypochondriasis				
Yes	37	57	0.633	0.426
No	900	1,645		
<hr/>				
Substance Misuse				
Yes	165	231	7.673	0.006
No	776	1,477		

\*n's may not add up to 941 because of missing observations (<1%)

\*\*n's may not add up to 1708 because of missing observations (<1%)

## **Appendix 2. Description of the Depressive Disorders Module in the Composite International Diagnostic Interview (CIDI), Spanish version 2.1**

This module begins with two screening questions concerning lifetime depression episodes. The first question asks whether the participant has ever felt sad, empty, or depressed for two weeks or longer. The second question asks whether the participant has ever lost interest in things he/she used to enjoy for a period of two weeks or longer. Participants who endorsed either question were given the full depression module. Following DSM-IV-TR diagnostic criteria, the full module asks questions whether the participant had ever felt the following symptoms for a period of two or more weeks: lack of energy, change in appetite, sleep problems, slowness/restlessness, feelings of worthlessness or guilt, lack of confidence, difficulty concentrating, and thoughts of death. To meet diagnostic criteria for MDD, five (or more) of the symptoms have been present during the same two-week period and represent a change from previous functioning; at least one of the symptoms is either: 1) depressed mood; or 2) loss of interest or pleasure (i.e., the screening questions). Symptoms are assessed for lifetime episodes, past year episodes, and during the most symptomatic episode (if applicable). The module also assesses about how many depressive episodes the individual has ever had, the age of onset and offset of symptoms, the severity of depressive symptoms, and whether they had sought help from a doctor during their depressive episodes (1).

- 1) World Health Organization. (1997) Composite International Diagnostic Interview (CIDI, Spanish Version 2.1). Geneva, Switzerland.