

# Topological data analysis distinguishes parameter regimes in the Anderson-Chaplain model of angiogenesis

John T. Nardini<sup>1</sup>, Bernadette J. Stolz<sup>2</sup>, Kevin B. Flores<sup>1</sup>, Heather A. Harrington<sup>2</sup>, Helen M. Byrne<sup>\*2</sup>

**1** Department of Mathematics, North Carolina State University, Raleigh, North Carolina, USA

**2** Mathematical Institute, University of Oxford, Oxford, OX2 6GG, UK

\* [helen.byrne@maths.ox.ac.uk](mailto:helen.byrne@maths.ox.ac.uk)

Parameter	baseline value
$D$	0.00035
$\chi$	0.38
$\rho$	0.34
$\beta$	0.05
$\gamma$	0.1
$\epsilon_1$	0.45
$\epsilon_2$	0.45
$k$	0.75

**S2 Table. Anderson-Chaplain model nondimensionalized parameters.**  
 Baseline nondimensionalized mechanistic parameters for the Chaplain-Anderson Model from [1].

## References

1. Anderson ARA, Chaplain MAJ. Continuous and Discrete Mathematical Models of Tumor-induced Angiogenesis. *Bulletin of Mathematical Biology*. 1998;60(5):857–899. doi:10.1006/bulm.1998.0042.