

Table E1: Proportion of current food allergies in three food groups among Black and White children with food allergy			
	Total	White	Black
	n(%)	n(%)	n (%)
Total	664 (100)	425 (100)	239 (100)
Type of Current Food Allergy			
Fin fish (>=1)	92 (13.9)	35 (8.2)	57 (23.9)
Salmon	65 (9.8)	25 (5.9)	40 (16.7)
Tuna	55 (8.3)	19 (4.5)	36 (15.1)
Tilapia	64 (9.6)	22 (5.2)	42 (17.6)
Trout	44 (6.6)	13 (3.1)	31 (12.9)
Halibut	49 (7.4)	16 (3.8)	33 (13.8)
Cod	62 (9.3)	21 (4.9)	41 (17.2)
Catfish	54 (8.1)	12 (2.8)	42 (17.6)
Swordfish	44 (6.6)	13 (3.1)	31 (12.9)
Grouper	43 (6.5)	12 (2.8)	31 (12.9)
Shellfish (>=1)	110 (16.6)	34 (8.0)	76 (31.8)
Shrimp	98 (14.8)	29 (6.8)	69 (28.9)
Lobster	77 (11.6)	22 (5.2)	55 (23.0)
Crab	79 (11.9)	23 (5.4)	56 (23.4)
Crayfish	63 (9.5)	13 (3.1)	50 (20.9)
Scallop	67 (10.1)	15 (3.5)	52 (21.8)
Oyster	62 (9.3)	15 (3.5)	47 (19.7)
Clam	61 (9.2)	15 (3.5)	46 (19.3)
Mussel	59 (8.9)	14 (3.3)	45 (18.8)
Tree nuts (>=1)	353 (53.2)	218 (51.3)	135 (56.5)
Almond	210 (31.6)	121(28.5)	89 (37.2)
Hazelnut	216 (32.5)	130 (30.6)	86 (35.9)
Cashew	290 (43.7)	179 (42.1)	111 (46.4)
Brazil	187 (28.2)	106 (24.9)	81 (33.9)
Macadamia	184 (27.7)	107 (25.2)	77 (32.2)
Pistachio	249 (37.5)	159 (37.4)	90 (37.7)
Pecan	222 (33.4)	131 (30.8)	91 (38.1)
Walnut	244 (36.8)	143 (33.7)	101 (42.3)

There were significant racial differences in the unadjusted analyses for all finfish (ie, salmon,

tuna, tilapia, trout, halibut, cod,catfish,swordfish and Grouper), all shellfish (ie, shrimp, lobster, crab, crayfish, scallop, oyster, clam, mussel) and three tree nuts (almond, brazil nuts, walnut). Due to the small numbers in each group an adjusted analysis was no done.

Journal Pre-proof