

Supplemental Online Content

Goldstick JE, Guy GP, Losby JL, Baldwin G, Myers M, Bohnert ASB. Changes in initial opioid prescribing practices after the 2016 release of the CDC Guideline for Prescribing Opioids for Chronic Pain. *JAMA Netw Open*. 2021;4(7):e2116860. doi:10.1001/jamanetworkopen.2021.16860

eFigure 1. Schematic Diagram of the Time Windows for Each Preguideline and Postguideline Cohort

eFigure 2. Proportion of All Beneficiaries Who Disenrolled From Coverage During Each Month

eFigure 3. Boxplot of Initial Prescription Durations Among Patients Who Were Opioid Naive in Each Cohort

eTable 1. Cancer and Palliative Care Diagnosis Codes Used for Exclusion Criteria

eTable 2. Age Distribution by Cohort Among Commercially Insured, Medicare Advantage Patients, and the Total Cohort

eTable 3. Race/Ethnicity Distribution by Cohort Among Commercially Insured, Medicare Advantage Patients, and the Total Cohort

eTable 4. Region Distribution by Cohort and Stratified by Insurer Type

eTable 5. Sex Distribution by Cohort Among Commercially Insured, Medicare Advantage Patients, and the Total Cohort

eTable 6. NHIS-Based Estimates of Commercially Insured Patient Demographics and Medicare Plus Insured Patient Demographics


eTable 7. Percentage of Enrollees Who Received a New Opioid Prescription During the Follow-up Period in Each Cohort

eTable 8. Mean Duration of Initial Opioid Prescription Among Patients Who Were Opioid Naive and Who Subsequently Received an Opioid Prescription by Cohort and Demographic; United States, 2011-2017

eTable 9. Percentage of Patients Who Were Opioid Naive and Who Subsequently Received an Opioid Prescription at 50 Morphine Milligram Equivalents Per Day or More


This supplemental material has been provided by the authors to give readers additional information about their work.

eFigure 1. Schematic Diagram of the Time Windows for Each Preguideline and Postguideline Cohort




Note: To qualify for each cohort individuals were required to: 1) be continuously enrolled for the entire 21-month period; 2) have 0 total Morphine Milligram Equivalents (MMEs) during their baseline period; and 3) Not having any diagnoses for cancer, palliative care, or end-of-life care at any time from 2011-2018.

eFigure 2. Proportion of All Beneficiaries Who Disenrolled From Coverage During Each Month



eFigure 3. Boxplot of Initial Prescription Durations Among Patients Who Were Opioid Naive in Each Cohort



Note: Whiskers extend up to five times the interquartile range above the 75th percentile

eTable 1. Cancer and Palliative Care Diagnosis Codes Used for Exclusion Criteria

Table X ICD-9 CM, ICD-10 CM, Cancer/Palliative Care Code Exclusions		
	ICD-9	ICD-10
Cancer (Malignant Neoplasm) Diagnosis*	140.X–208.X, 209.0–209.3, V10.X	C00X–C97X
Palliative Care	V66.7	Z51.5
*Excluding benign neoplasms or neoplasms of uncertain behavior		

eTable 2. Age Distribution by Cohort Among Commercially Insured, Medicare Advantage Patients, and the Total Cohort

Commercial				
Year	Age 18-34	Age 35-54	Age 55-74	Age 75+
Pre-Guideline Cohort 1	1,171,553 (32.0%)	1,737,007 (47.4%)	713,343 (19.5%)	42,275 (1.2%)
Pre-Guideline Cohort 2	1,193,784 (32.5%)	1,721,370 (46.9%)	723,113 (19.7%)	30,174 (0.8%)
Pre-Guideline Cohort 3	1,104,662 (33.5%)	1,534,174 (46.5%)	632,629 (19.2%)	26,162 (0.8%)
Pre-Guideline Cohort 4	1,182,535 (33.9%)	1,605,676 (46.1%)	670,712 (19.2%)	25,336 (0.7%)
Post-Guideline Cohort 1	1,233,130 (33.5%)	1,675,721 (45.6%)	742,793 (20.2%)	26,060 (0.7%)
Post-Guideline Cohort 2	1,328,032 (33.8%)	1,776,120 (45.2%)	795,678 (20.3%)	29,371 (0.7%)
Medicare				
Year	Age 18-34	Age 35-54	Age 55-74	Age 75+
Pre-Guideline Cohort 1	3,185 (0.3%)	29,354 (2.8%)	602,388 (57.8%)	406,988 (39.1%)
Pre-Guideline Cohort 2	3,754 (0.3%)	33,285 (2.8%)	699,762 (58.2%)	464,570 (38.7%)
Pre-Guideline Cohort 3	3,998 (0.3%)	33,736 (2.9%)	684,874 (58.1%)	455,483 (38.7%)
Pre-Guideline Cohort 4	4,477 (0.3%)	34,275 (2.6%)	758,808 (57.7%)	517,991 (39.4%)
Post-Guideline Cohort 1	5,130 (0.3%)	37,450 (2.4%)	891,168 (57.2%)	623,506 (40.0%)
Post-Guideline Cohort 2	6,842 (0.4%)	49,002 (2.6%)	1,097,259 (57.6%)	751,784 (39.5%)
Total				
Year	Age 18-34	Age 35-54	Age 55-74	Age 75+
Pre-Guideline Cohort 1	1,174,738 (25.0%)	1,766,361 (37.5%)	1,315,731 (28.0%)	449,263 (9.5%)
Pre-Guideline Cohort 2	1,197,538 (24.6%)	1,754,655 (36.0%)	1,422,875 (29.2%)	494,744 (10.2%)
Pre-Guideline Cohort 3	1,108,660 (24.8%)	1,567,910 (35.0%)	1,317,503 (29.4%)	481,645 (10.8%)
Pre-Guideline Cohort 4	1,187,012 (24.7%)	1,639,951 (34.2%)	1,429,520 (29.8%)	543,327 (11.3%)
Post-Guideline Cohort 1	1,238,260 (23.7%)	1,713,171 (32.7%)	1,633,961 (31.2%)	649,566 (12.4%)
Post-Guideline Cohort 2	1,334,874 (22.9%)	1,825,122 (31.3%)	1,892,937 (32.4%)	781,155 (13.4%)

eTable 3. Race/Ethnicity Distribution by Cohort Among Commercially Insured, Medicare Advantage Patients, and the Total Cohort

Commercial					
Year	Unknown	Asian	Black	Hispanic	White
Pre-Guideline Cohort 1	243,323 (6.6%)	199,454 (5.4%)	345,489 (9.4%)	396,516 (10.8%)	2,479,396 (67.7%)
Pre-Guideline Cohort 2	241,506 (6.6%)	208,108 (5.7%)	340,178 (9.3%)	401,333 (10.9%)	2,477,316 (67.5%)
Pre-Guideline Cohort 3	219,272 (6.6%)	196,918 (6.0%)	271,081 (8.2%)	395,276 (12.0%)	2,215,080 (67.2%)
Pre-Guideline Cohort 4	238,780 (6.9%)	207,589 (6.0%)	291,130 (8.4%)	431,157 (12.4%)	2,315,603 (66.5%)
Post-Guideline Cohort 1	240,044 (6.5%)	221,202 (6.0%)	308,458 (8.4%)	469,137 (12.8%)	2,438,863 (66.3%)
Post-Guideline Cohort 2	413,438 (10.5%)	224,491 (5.7%)	310,124 (7.9%)	488,546 (12.4%)	2,492,602 (63.4%)
Medicare					
Year	Unknown	Asian	Black	Hispanic	White
Pre-Guideline Cohort 1	179,412 (17.2%)	31,210 (3.0%)	87,398 (8.4%)	93,530 (9.0%)	650,365 (62.4%)
Pre-Guideline Cohort 2	183,748 (15.3%)	50,485 (4.2%)	105,652 (8.8%)	111,526 (9.3%)	749,960 (62.4%)
Pre-Guideline Cohort 3	180,123 (15.3%)	53,078 (4.5%)	95,674 (8.1%)	116,198 (9.9%)	733,018 (62.2%)
Pre-Guideline Cohort 4	196,927 (15.0%)	57,703 (4.4%)	107,113 (8.1%)	144,247 (11.0%)	809,561 (61.5%)
Post-Guideline Cohort 1	237,734 (15.3%)	64,285 (4.1%)	133,992 (8.6%)	159,567 (10.2%)	961,676 (61.8%)
Post-Guideline Cohort 2	289,295 (15.2%)	69,534 (3.7%)	191,135 (10.0%)	187,376 (9.8%)	1,167,547 (61.3%)
Total					
Year	Unknown	Asian	Black	Hispanic	White
Pre-Guideline Cohort 1	422,735 (9.0%)	230,664 (4.9%)	432,887 (9.2%)	490,046 (10.4%)	3,129,761 (66.5%)
Pre-Guideline Cohort 2	425,254 (8.7%)	258,593 (5.3%)	445,830 (9.2%)	512,859 (10.5%)	3,227,276 (66.3%)
Pre-Guideline Cohort 3	399,395 (8.9%)	249,996 (5.6%)	366,755 (8.2%)	511,474 (11.4%)	2,948,098 (65.9%)
Pre-Guideline Cohort 4	435,707 (9.1%)	265,292 (5.5%)	398,243 (8.3%)	575,404 (12.0%)	3,125,164 (65.1%)
Post-Guideline Cohort 1	477,778 (9.1%)	285,487 (5.5%)	442,450 (8.5%)	628,704 (12.0%)	3,400,539 (65.0%)
Post-Guideline Cohort 2	702,733 (12.0%)	294,025 (5.0%)	501,259 (8.6%)	675,922 (11.6%)	3,660,149 (62.7%)

eTable 4. Region Distribution by Cohort and Stratified by Insurer Type

Commercial						
Year	West	Midwest	South	Northeast	Pacific	Unknown
Pre-Guideline Cohort 1	730,598 (19.9%)	944,594 (25.8%)	1,590,444 (43.4%)	380,466 (10.4%)	2,481 (0.1%)	15,595 (0.4%)
Pre-Guideline Cohort 2	750,380 (20.5%)	964,323 (26.3%)	1,555,238 (42.4%)	368,040 (10.0%)	2,510 (0.1%)	27,950 (0.8%)
Pre-Guideline Cohort 3	719,034 (21.8%)	911,726 (27.6%)	1,297,445 (39.3%)	330,208 (10.0%)	2,223 (0.1%)	36,991 (1.1%)
Pre-Guideline Cohort 4	748,507 (21.5%)	958,751 (27.5%)	1,374,387 (39.4%)	348,808 (10.0%)	2,666 (0.1%)	51,140 (1.5%)
Post-Guideline Cohort 1	784,533 (21.3%)	1,008,291 (27.4%)	1,460,199 (39.7%)	377,425 (10.3%)	2,831 (0.1%)	44,425 (1.2%)
Post-Guideline Cohort 2	838,609 (21.3%)	1,039,149 (26.4%)	1,495,979 (38.1%)	379,416 (9.7%)	2,852 (0.1%)	173,196 (4.4%)
Medicare						
Year	West	Midwest	South	Northeast	Pacific	Unknown
Pre-Guideline Cohort 1	375,379 (36.0%)	215,979 (20.7%)	338,290 (32.5%)	85,827 (8.2%)	9,402 (0.9%)	17,038 (1.6%)
Pre-Guideline Cohort 2	383,575 (31.9%)	247,390 (20.6%)	373,364 (31.1%)	170,918 (14.2%)	9,779 (0.8%)	16,345 (1.4%)
Pre-Guideline Cohort 3	382,503 (32.5%)	262,451 (22.3%)	324,018 (27.5%)	180,257 (15.3%)	8,451 (0.7%)	20,411 (1.7%)
Pre-Guideline Cohort 4	397,713 (30.2%)	298,967 (22.7%)	398,725 (30.3%)	206,748 (15.7%)	8,966 (0.7%)	4,432 (0.3%)
Post-Guideline Cohort 1	431,229 (27.7%)	343,839 (22.1%)	524,222 (33.7%)	241,461 (15.5%)	11,634 (0.7%)	4,869 (0.3%)
Post-Guideline Cohort 2	483,479 (25.4%)	431,406 (22.6%)	693,799 (36.4%)	284,398 (14.9%)	6,461 (0.3%)	5,344 (0.3%)
Total						
Year	West	Midwest	South	Northeast	Pacific	Unknown
Pre-Guideline Cohort 1	1,105,977 (23.5%)	1,160,573 (24.7%)	1,928,734 (41.0%)	466,293 (9.9%)	11,883 (0.3%)	32,633 (0.7%)
Pre-Guideline Cohort 2	1,133,955 (23.3%)	1,211,713 (24.9%)	1,928,602 (39.6%)	538,958 (11.1%)	12,289 (0.3%)	44,295 (0.9%)
Pre-Guideline Cohort 3	1,101,537 (24.6%)	1,174,177 (26.2%)	1,621,463 (36.2%)	510,465 (11.4%)	10,674 (0.2%)	57,402 (1.3%)
Pre-Guideline Cohort 4	1,146,220 (23.9%)	1,257,718 (26.2%)	1,773,112 (36.9%)	555,556 (11.6%)	11,632 (0.2%)	55,572 (1.2%)
Post-Guideline Cohort 1	1,215,762 (23.2%)	1,352,130 (25.8%)	1,984,421 (37.9%)	618,886 (11.8%)	14,465 (0.3%)	49,294 (0.9%)
Post-Guideline Cohort 2	1,322,088 (22.7%)	1,470,555 (25.2%)	2,189,778 (37.5%)	663,814 (11.4%)	9,313 (0.2%)	178,540 (3.1%)

Regional categories based on census categories West, Midwest, Northeast, and South (https://www2.census.gov/geo/pdfs/maps-data/maps/reference/us_regdiv.pdf). Additionally, added “Pacific” to refer to Puerto Rico, Hawaii, and Alaska.

eTable 5. Sex Distribution by Cohort Among Commercially Insured, Medicare Advantage Patients, and the Total Cohort

Commercial			
Year	Male	Female	Unknown
Pre-Guideline Cohort 1	1,876,731	1,786,748	699
	(51.2%)	(48.8%)	(0.0%)
Pre-Guideline Cohort 2	1,888,038	1,779,763	640
	(51.5%)	(48.5%)	(0.0%)
Pre-Guideline Cohort 3	1,727,385	1,569,691	551
	(52.4%)	(47.6%)	(0.0%)
Pre-Guideline Cohort 4	1,825,382	1,658,428	449
	(52.4%)	(47.6%)	(0.0%)
Post-Guideline Cohort 1	1,922,439	1,754,910	355
	(52.3%)	(47.7%)	(0.0%)
Post-Guideline Cohort 2	2,071,439	1,857,466	296
	(52.7%)	(47.3%)	(0.0%)
Medicare			
Year	Male	Female	Unknown
Pre-Guideline Cohort 1	437,213	604,702	0
	(42.0%)	(58.0%)	(0.0%)
Pre-Guideline Cohort 2	508,666	692,705	0
	(42.3%)	(57.7%)	(0.0%)
Pre-Guideline Cohort 3	503,385	674,706	0
	(42.7%)	(57.3%)	(0.0%)
Pre-Guideline Cohort 4	551,168	764,383	0
	(41.9%)	(58.1%)	(0.0%)
Post-Guideline Cohort 1	643,805	913,449	0
	(41.3%)	(58.7%)	(0.0%)
Post-Guideline Cohort 2	788,664	1,116,223	0
	(41.4%)	(58.6%)	(0.0%)
Total			
Year	Male	Female	Unknown
Pre-Guideline Cohort 1	2,313,944	2,391,450	699
	(49.2%)	(50.8%)	(0.0%)
Pre-Guideline Cohort 2	2,396,704	2,472,468	640
	(49.2%)	(50.8%)	(0.0%)
Pre-Guideline Cohort 3	2,230,770	2,244,397	551
	(49.8%)	(50.1%)	(0.0%)
Post-Guideline Cohort 4	2,376,550	2,422,811	449
	(49.5%)	(50.5%)	(0.0%)
Post-Guideline Cohort 1	2,566,244	2,668,359	355
	(49.0%)	(51.0%)	(0.0%)
Pre-Guideline Cohort 2	2,860,103	2,973,689	296
	(49.0%)	(51.0%)	(0.0%)

eTable 6. NHIS-Based Estimates of Commercially Insured Patient Demographics and Medicare Plus Insured Patient Demographics

Among All Commercially Insured				
Year	Age (mean; 95% CI)	Gender (% male, 95% CI)	Race (%White, 95% CI)	Race(%Black, 95% CI)
2012	38.24 (38.06-38.41)	49.0 (48.6-49.4)	83.4 (82.7-84.0)	9.7 (9.1-10.2)
2013	38.29 (38.11-38.46)	49.2 (48.8-49.5)	83.0 (82.3-83.7)	9.8 (9.2-10.3)
2014	38.39 (38.23-38.56)	49.3 (49.0-49.7)	82.7 (82.0-83.3)	9.8 (9.2-10.3)
2015	38.19 (38.02-38.36)	49.1 (48.8-49.5)	81.9 (81.2-82.6)	10.2 (9.7-10.8)
2016	38.38 (38.21-38.55)	49.4 (49.0-49.8)	81.3 (80.2-82.5)	10.4 (9.6-11.3)
2017	38.37 (38.18-38.56)	49.3 (48.9-49.7)	81.1 (79.7-82.4)	10.4 (9.5-11.4)
Among those Enrolled in Medicare Plus Choice				
Year	Age (mean; 95% CI)	Gender(% male; 95% CI)	Race (%White; 95% CI)	Race(%Black; 95% CI)
2012	72.21 (71.88-72.54)	43.1 (41.3-44.9)	86.5 (84.6-88.2)	9.4 (8.0-11.1)
2013	71.52 (71.17-71.86)	42.7 (41.1-44.2)	86.6 (84.9-88.1)	8.8 (7.6-10.1)
2014	72.25 (71.93-72.56)	43.9 (42.3-45.6)	86.8 (85.2-88.3)	8.7 (7.5-10.1)
2015	72.11 (71.82-72.40)	44.4 (42.9-45.9)	84.9 (83.1-86.6)	10.1 (8.8-11.6)
2016	71.88 (71.59-72.17)	43.7 (42.3-45.2)	84.8 (82.8-86.6)	10 (8.7-11.6)
2017	71.9 (71.61-72.20)	44.1 (42.7-45.6)	84.2 (82.1-86.1)	10.1 (8.5-11.9)

Note: For comparison with the cohort composition, 2012, 2013, 2014, 2015, 2016, 2017 are most closely aligned temporally with Pre-Guideline Cohort 1, Pre-Guideline Cohort 2, Pre-Guideline Cohort 3, Pre-Guideline Cohort 4, Post-Guideline Cohort 1, and Post-Guideline Cohort 2, respectively.

eTable 7. Percentage of Enrollees Who Received a New Opioid Prescription During the Follow-up Period in Each Cohort

	Total	Female	Male	White	Non-white	Commercial	Medicare Advantage
Pre-Guideline 1	558,175 (11.9%)	310,650 (13.0%)	247,425 (10.7%)	385,855 (12.3%)	134,612 (11.7%)	441,490 (12.0%)	116,685 (11.1%)
Pre-Guideline 2	556,234 (11.4%)	309,423 (12.5%)	246,732 (10.3%)	383,397 (11.9%)	136,385 (11.2%)	421,652 (11.4%)	134,582 (11.2%)
Pre-Guideline 3	472,511 (10.6%)	259,200 (11.5%)	213,251 (9.6%)	325,781 (11.1%)	114,895 (10.2%)	348,682 (10.5%)	123,829 (10.5%)
Pre-Guideline 4	487,131 (10.1%)	269,788 (11.1%)	217,277 (9.1%)	327,349 (10.5%)	120,133 (9.7%)	345,900 (9.8%)	141,231 (10.7%)
Post-Guideline 1	505,155 (9.6%)	280,818 (10.5%)	224,310 (8.7%)	338,484 (10.0%)	127,004 (9.4%)	352,350 (9.5%)	152,805 (9.8%)
Post-Guideline 2	532,962 (9.1%)	297,391 (10.0%)	235,554 (8.2%)	349,555 (9.6%)	134,358 (9.1%)	343,514 (8.7%)	189,448 (9.9%)
(actual-predicted)	+0.1%	+0.1%	+0.1%	+0.2%	+0.7%	+0.1%	-0.1%

Note: The final row shows the difference between the observed level in the Post-Guideline group 2, and what was projected based on a linear extrapolation of the pre-guideline trend.

Note 2: Those with missing values for race were excluded from the “white” and “non-white” groups, but were not excluded from any other columns

eTable 8. Mean Duration of Initial Opioid Prescription Among Patients Who Were Opioid Naive and Who Subsequently Received an Opioid Prescription by Cohort and Demographic; United States, 2011-2017

	Overall	Female	Male	White	Non-white	Commercial Insurance	Medicare Advantage
Pre-Guideline 1	6.17	6.31	6.00	6.01	6.54	5.50	8.73
Pre-Guideline 2	6.40	6.57	6.19	6.22	6.79	5.61	8.87
Pre-Guideline 3	6.49	6.65	6.30	6.29	6.93	5.62	8.93
Pre-Guideline 4	6.64	6.84	6.39	6.35	7.04	5.62	9.14
Post-Guideline 1	6.49	6.70	6.24	6.25	6.93	5.52	8.72
Post-Guideline 2	6.38	6.61	6.09	6.11	6.89	5.22	8.49
(actual-pred)	-0.57	-0.56	-0.59	-0.50	-0.51	-0.51	-0.88
% change (difference/predicted)	-8.2%	-7.8%	-8.8%	-7.5%	-6.9%	-8.8%	-9.4%

Note: The final row shows the difference between the observed level in the Post-Guideline group 2, and what was projected based on a linear extrapolation of the pre-guideline trend. The % reduction (diff/pred) entry shows the percent change in the actual level, relative to the projection.

Note 2: Those with missing values for race were excluded from the “Non-white” group, but were not excluded from any other columns.

eTable 9. Percentage of Patients Who Were Opioid Naive and Who Subsequently Received an Opioid Prescription at 50 Morphine Milligram Equivalents Per Day or More

	Total	Female	Male	White	Non-white	Commercial	Medicare Advantage
Pre-Guideline 1	130,883 (23.4%)	71,046 (22.9%)	59,819 (24.2%)	95,532 (24.8%)	26,672 (19.8%)	103,160 (23.4%)	27,723 (23.8%)
Pre-Guideline 2	120,861 (21.7%)	65,172 (21.1%)	55,675 (22.6%)	88,488 (23.1%)	24,395 (17.9%)	94,390 (22.4%)	26,471 (19.7%)
Pre-Guideline 3	110,017 (23.3%)	58,045 (22.4%)	51,959 (24.4%)	81,103 (24.9%)	21,661 (18.9%)	86,920 (24.9%)	23,097 (18.7%)
Pre-Guideline 4	110,734 (22.7%)	59,049 (21.9%)	51,672 (23.8%)	80,912 (24.7%)	21,375 (17.8%)	84,359 (24.4%)	26,375 (18.7%)
Post-Guideline 1	110,088 (21.8%)	58,693 (20.9%)	51,390 (22.9%)	80,598 (23.8%)	21,172 (16.7%)	82,703 (23.5%)	27,385 (17.9%)
Post-Guideline 2	112,312 (21.1%)	60,116 (20.2%)	52,189 (22.2%)	80,813 (23.1%)	21,591 (16.1%)	79,278 (23.1%)	33,034 (17.4%)
(actual-pred)	-1.5%	-1.2%	-1.7%	-1.8%	-0.7%	-2.6%	+2.2%

Note: The final row shows the difference between the observed level in the Post-Guideline group 2, and what was projected based on a linear extrapolation of the pre-guideline trend.

Note 2: Those with missing values for race were excluded from the “Non-white” group, but were not excluded from any other columns.