

**Serum 8-isoprostane levels in patients with resistant oral lichen  
planus before and after treatment with lycopene: A  
Randomized Clinical Trial**

Aliaa Abdelmoniem Bedeir Eita\*, Azza Mohamed Zaki,  
Sabah Abdelhady Mahmoud

**Supplementary Table S1: Median, Inter Quartile Range (IQR) and inter-group comparisons of Escudier et al. lesion scores at all assessment times**

Site score	Lycopene group (n=10)	Corticosteroids group (n=10)	Mann Whitney U p value
	Median (IQR)		
<b>Baseline</b>	4.00 (2.75, 6.00)	3.00 (2.75, 4.50)	P = 0.25
<b>4 weeks</b>	4.00 (1.75, 4.50)	2.00 (1.00, 2.50)	P = 0.09
<b>8 weeks</b>	3.50 (1.75, 4.50)	2.00 (1.00, 2.00)	P = 0.09
Severity score	Median (IQR)		Mann Whitney U p value
<b>Baseline</b>	4.00 (3.00, 5.00)	5.90 (4.50, 6.75)	P = 0.08
<b>4 weeks</b>	2.50 (1.00, 5.00)	2.00 (1.00, 4.75)	P = 0.91
<b>8 weeks</b>	1.00 (0.00, 2.50)	2.00 (1.00, 3.25)	P = 0.28
Activity score	Median (IQR)		Mann Whitney U p value
<b>Baseline</b>	6.50 (5.00, 8.25)	8.00 (5.00, 10.00)	P = 0.63
<b>4 weeks</b>	4.50 (1.00, 7.25)	2.00 (1.00, 4.75)	P = 0.53
<b>8 weeks</b>	1.50 (0.00, 3.00)	2.00 (1.00, 3.25)	P = 0.44

\*statistically significant at p value < 0.05

n: number of patients