Table S1. Baseline study group characteristics in the total population and by adherence to diabetic retinopathy evaluation

	Total Population	Received Examination	Did Not Follow-up
Parameter	N	n	n
No. patients	1397	727	670
Age, years; mean (SD)	54.1 (13.9)	55.7 (14.0)	52.5 (13.7)
Gender Male Female  Race Asian or Pacific Islander Black or African American Hispanic or Latino	600 (42.9%)	315 (43.3%)	285 (42.5%)
	797 (57.1%)	412 (56.7%)	385 (57.5%)
	30 (2.1%)	22 (3.0%)	8 (1.1%)
	580 (41.5%)	293 (40.3%)	287 (42.8%)
	488 (34.9%)	228 (31.4%)	260 (38.8%)
Non-Hispanic White Other	239 (17.1%)	153 (21.0%)	86 (12.8%)
	60 (4.3%)	31 (4.3%)	29 (4.3%)
Insurance Status Commercial Medicaid Medicare Medicare Managed Self-Pay Other	105 (7.5%)	67 (9.2%)	38 (5.7%)
	642 (46.0%)	305 (42.0%)	337 (50.3%)
	385 (27.6%)	221 (30.4%)	164 (24.5%)
	154 (11.0%)	94 (12.9%)	60 (9.0%)
	107 (7.7%)	38 (5.2%)	69 (10.3%)
	4 (0.3%)	2 (0.3%)	2 (0.3%)
National ADI Quintile Quintile 1 (Most disadvantaged) Quintile 2 Quintile 3 Quintile 4 Quintile 5 (Least disadvantaged)	176 (12.6%)	69 (9.5%)	107 (16.0%)
	379 (27.1%)	189 (26.0%)	190 (28.4%)
	429 (30.7%)	237 (32.6%)	192 (28.7%)
	312 (22.3%)	175 (24.1%)	137 (20.4%)
	101 (7.2%)	57 (7.8%)	44 (6.6%)

ADI = Area Deprivation Index; SD = Standard deviation