

Isolation and characterization of novel bacteriophages as a potential therapeutic option for *Escherichia coli* urinary tract infections

Edgar González-Villalobos^{a,b,c,d}, Rosa María Ribas-Aparicio^a, Gerardo Erbey Rodea Montealegre^{b,c}, Laura Belmont-Monroy^{a,b,c}, Yerisaidy Ortega-García^{a,b,c}, Gerardo Aparicio-Ozores^a, José Luis Balcázar^{d,e}, Carlos Alberto Eslava-Campos^{b,c}, Ulises Hernández-Chiñas^{b,c}, José Molina-López^{b,c*}

^a Departamento de Microbiología, Escuela Nacional de Ciencias Biológicas (ENCB), Instituto Politécnico Nacional (IPN). Prolongación de Carpio y Plan de Ayala s/n, Colonia Santo Tomás, C.P. 11340, Mexico City, Mexico.

^b Unidad Periférica de Investigación Básica y Clínica en Enfermedades Infecciosas; Departamento de Salud Pública/División de Investigación, Facultad de Medicina, UNAM, C.P. 04510. Mexico City, Mexico.

^c Laboratorio de Patogenicidad Bacteriana, Unidad de Hemato-Oncología e Investigación. Hospital Infantil de México Federico Gómez/Facultad de Medicina UNAM, C.P. 06720 Mexico City, Mexico.

^d Catalan Institute for Water Research (ICRA). Girona 17003, Spain.

^e University of Girona. Girona 17004, Spain.

*Corresponding author:

Dr. José Molina-López

Phone: +52 55 52 28 99 17.

E-mail: joseml@unam.mx

Supplementary material

Supplementary Table S1. Mass spectrometry data from ϕ Ec3 and ϕ Ec4 phages. In both cases, similar proteins were detected by mass spectrometry. ID denotes the reference of the identified proteins. M/Z indicates the mass-charge ratio. Peptide refers to a protein sequence.

ID	Coverage	M/Z	Peptide
RNA2 polyprotein	6.5%	871.257	DMISDFK
		887.239	ISTSVAPGR
		1364.076	WITAGLVMPTFK
		1758.347	ISTSVAPGRAQWISER
		1792.164	MFSWKWGSGPIDNPEK
		2383.480	ITASADPVYTLSVPHWLIHKK
Endolysin	19.0%	862.116	QAEDMLR
		1046.608	SGDNLTKIAK
		1347.711	SGDTLSKIAVDNK
		2086.956	QISQAGINLIKSFEGLQLK

Supplementary Fig. S1. Analysis of ϕ Ec3 and ϕ Ec4 protein profiles (15% SDS-PAGE with silver staining). ϕ Ec3, two proteins of 37 kDa and 11 kDa were identified. ϕ Ec4, two proteins of 37 kDa and 13 kDa were identified.

