

Appendix 3. Stratum-specific frequency and percent linking to care and prevalence ratios and prevalence differences comparing intervention effect by sex, age, household food insecurity, and SSQ score

	Percent seeking care				Prevalence ratio	(95% CI)	Prevalence difference	(95% CI)	p-value	N
	Conditional incentive		No incentive							
	No.	%	No.	%						
Sex										
Men	338	23.9	364	26.9	0.93	(0.81, 1.07)	-0.018	(-0.051, 0.015)	0.295	2,767
Women	724	31.7	711	33.94	0.95	(0.87, 1.05)	-0.015	(-0.045, 0.015)	0.334	4,379
Age *										
Younger adults (up to 25 years)	322	31.05	320	33.61	0.96	(0.84, 1.11)	-0.011	(-0.052, 0.030)	0.593	1,989
Older adults (26 years and older)	740	27.86	755	30.3	0.94	(0.86, 1.04)	-0.017	(-0.045, 0.011)	0.225	5,148
Food insecurity										
Food-secure household	412	24.74	423	27.94	0.9	(0.80, 1.03)	-0.025	(-0.058, 0.007)	0.123	3,179
Food-insecure household	650	31.97	652	33.71	0.98	(0.89, 1.08)	-0.007	(-0.039, 0.025)	0.667	3,967
SSQ										
Lower SSQ (0-8)	476	26.4	422	27.17	0.95	(0.87, 1.04)	-0.003	(-0.033, 0.027)	0.833	3,356
Higher SSQ (9-14)	586	30.92	653	34.46	0.95	(0.88, 1.03)	-0.03	(-0.064, 0.005)	0.092	3,790

SSQ = Shona Symptom Questionnaire. Subgroup analyses of sex and age adjusted for educational attainment, household food insecurity, and SSQ. Subgroup analysis of food insecurity adjusted for educational attainment and SSQ. Subgroup analysis of SSQ adjusted for educational attainment and food insecurity.

* 9 Respondents missing age