

Supplementary table

Supplemental Table 1: Receiver operating characteristics (ROC) for clinical scores and selected laboratory tests in relation to 30-day mortality for patients with suspected sepsis.

	AUC	N
Charlson comorbidity index	0.65 (0.62-0.68)	1227
SOFA	0.73 (0.71-0.76)	1229
qSOFA	0.67 (0.65-0.70)	1231
NRBC	0.60 (0.57-0.63)	1231
Platelets	0.56 (0.53-0.59)	1228
Bilirubin	0.54 (0.51-0.57)	1226
Glasgow coma scale	0.67 (0.64-0.69)	1231
Creatinine	0.62 (0.59-0.65)	1231
Albumin	0.66 (0.63-0.69)	1226
CRP	0.55 (0.52-0.58)	1231
D-dimer	0.66 (0.64-0.69)	1133
Lactate	0.68 (0.65-0.70)	1072
Lactate dehydrogenase	0.68 (0.65-0.72)	713
Leukocytes	0.54 (0.51-0.57)	1231
Lymphocytes	0.50 (0.48-0.53)	1194
Neutrophils	0.52 (0.50-0.52)	1194
Neutrophil/lymphocyte ratio	0.50 (0.48-0.53)	1194

AUC=Area under the curve, with 95 % confidence intervals.

Supplementary Figure legend

Supplemental Figure 1: Relationship between NRBC score and other predictive variables for patients with suspected sepsis. Charlson score (A), SOFA score (B), Glasgow coma scale (GCS) (C) and laboratory results (D-J) grouped according to NRBC results. Red lines represent the median value.

