

Supplementary Table SII Number and percentage of offspring diabetes in birth cohort 1996–2014. Values are n (per 100 000).

Groups	Type 1 diabetes	Type 2 diabetes	Nutrition-related diabetes	Other specified diabetes	Other unspecified diabetes	Total
No maternal PCOS/anovulatory infertility						
First diagnosis	8934 (832.6)	231 (21.5)	7 (0.7)	84 (7.8)	135 (12.6)	9391 (875.2)
Second diagnosis	8716 (812.2)	340 (31.7)	9 (0.8)	124 (11.6)	202 (18.8)	9391 (875.2)
Maternal PCOS/anovulatory infertility						
First diagnosis	192 (777.9)	9 (36.5)	0 (0)	7 (28.4)	6 (24.3)	214 (867)
Second diagnosis	197 (798.2)	11 (44.6)	0 (0)	1 (4.1)	5 (20.3)	214 (867)

PCOS (polycystic ovary syndrome)/anovulatory infertility was identified by ICD-9 256.4, 628.0, ICD-10 E28.2, and N97.0. Diabetes was identified based on ICD codes, E10 for Type 1 diabetes, E11 for Type 2 diabetes, E12 for nutrition-related diabetes, E13 for other specified diabetes, and E14 for other unspecified diabetes.